


Study of 52,000 men uncovers the genetics underlying male pattern baldness

February 14 2017



Genetics of male pattern baldness are shown. Credit: Douglas Robertson, University of Edinburgh Centre for Cognitive Ageing and Cognitive Epidemiology

A genomic study of baldness identified more than 200 genetic regions involved in this common but potentially embarrassing condition. These genetic variants could be used to predict a man's chance of severe hair loss. The study, led by Saskia Hagenaars and W. David Hill of The University of Edinburgh, United Kingdom, is published February 14th, 2017 in *PLOS Genetics*.

Before this new study, only a handful of genes related to [baldness](#) had

been identified. The University of Edinburgh scientists examined genomic and health data from over 52,000 male participants of the UK Biobank, performing a genome-wide association study of baldness. They pinpointed 287 [genetic regions](#) linked to the condition. The researchers created a formula to try and predict the chance that a person will go bald, based on the presence or absence of certain [genetic](#) markers. Accurate predictions for an individual are still some way off, but the results can help to identify sub-groups of the population for which the risk of hair loss is much higher.

The study is the largest genetic analysis of male pattern baldness to date. Many of the identified genes are related to hair structure and development. They could provide possible targets for drug development to treat baldness or related conditions.

Saskia Hagenaars, a PhD student from The University of Edinburgh's Centre for Cognitive Ageing and Cognitive Epidemiology, who jointly led the research, said: "We identified hundreds of new genetic signals. It was interesting to find that many of the genetics signals for male pattern baldness came from the X chromosome, which men inherit from their mothers."

Dr David Hill, who co-led the research, said: "In this study, data were collected on hair loss pattern but not age of onset; we would expect to see an even stronger genetic signal if we were able to identify those with early-onset hair loss."

The study's principal investigator, Dr Riccardo Marioni, from The University of Edinburgh's Centre for Genomic and Experimental Medicine, said: "We are still a long way from making an accurate prediction for an individual's hair loss pattern. However, these results take us one step closer. The findings pave the way for an improved understanding of the genetic causes of [hair loss](#)."

More information: Hageaars SP, Hill WD, Harris SE, Ritchie SJ, Davies G, Liewald DC, et al. (2017) Genetic prediction of male pattern baldness. *PLoS Genet* 13(2): e1006594. [DOI: 10.1371/journal.pgen.1006594](https://doi.org/10.1371/journal.pgen.1006594)

Provided by Public Library of Science

Citation: Study of 52,000 men uncovers the genetics underlying male pattern baldness (2017, February 14) retrieved 23 April 2024 from <https://medicalxpress.com/news/2017-02-men-uncovers-genetics-underlying-male.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.