

Modern housing associated with reduced malaria risk in sub-Saharan Africa

February 21 2017

Modern houses—with metal roofs and finished walls—are associated with a more than 9 percent reduction in the odds of malaria in children in sub-Saharan Africa when compared to more traditional thatched houses, according to a study published in *PLOS Medicine* by Lucy Tusting of the University of Oxford, UK, and colleagues at the London School of Hygiene & Tropical Medicine, UK, Durham University, UK, and the University of Southampton, UK.

Insecticide-treated bednets and house spraying have been effective in reducing the prevalence of [malaria](#) since the turn of the 21st century, but other approaches are needed for sustainable elimination of the mosquito-transmitted, parasitic disease. Some evidence has suggested that modern houses may protect against the parasite but few studies have rigorously evaluated the association between improved housing and malaria risk. In the new work, researchers analyzed data on malaria prevalence and housing using data collected in 29 surveys carried out in 21 African countries between 2008 and 2015. Information on malaria status—as tested using a blood smear or rapid test—was available for 139,318 children under age 5 living in 84,153 households.

The proportion of children with malaria detectable in their blood varied by survey, ranging from 0.4% to 45.5% among children living in modern houses and from 0.4% to 70.6% among children living in traditional homes. Across all surveys, modern housing was associated with a 9% to 14% reduction in the odds of [malaria infection](#), after controlling for household wealth and use of insecticides. By comparison, [children](#)

sleeping under insecticide-treated bednets had a 15% to 16% reduction in odds of testing positive for the disease.

Lead author Lucy Tusting from the University of Oxford, said: "Good housing is a core pillar of public health, but is not used widely for malaria control. Well-built housing can block mosquitoes from entering homes and prevent them from transmitting malaria to the people who live there.

"This is the first study to compare housing and insecticide-treated nets for [malaria control](#) across a range of countries in sub-Saharan Africa. Our study suggests good housing could be an important tool in tackling malaria. This is a welcome finding at a time when we are facing increasing resistance to our most effective insecticides and drugs. We now need to assess the impact of housing improvements on malaria in field trials and to work with architects and urban planners to incorporate protective designs into housing in regions at risk of malaria."

The authors note that the effectiveness of improving [housing](#) will vary depending on the location. While many mosquitoes enter homes to bite humans at night, outdoor malaria transmission is more common in some places, meaning interventions centered on the home will have less impact.

More information: Tusting LS, Bottomley C, Gibson H, Kleinschmidt I, Tatem AJ, Lindsay SW, et al. (2017) Housing Improvements and Malaria Risk in Sub-Saharan Africa: A Multi-Country Analysis of Survey Data. *PLoS Med* 14(2): e1002234. [DOI: 10.1371/journal.pmed.1002234](#)

Provided by Public Library of Science

Citation: Modern housing associated with reduced malaria risk in sub-Saharan Africa (2017, February 21) retrieved 11 July 2024 from <https://medicalxpress.com/news/2017-02-modern-housing-malaria-sub-saharan-africa.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.