

More patients with early-stage breast cancer may be able to avoid chemotherapy in the future

February 15 2017

Women with early-stage breast cancer who had an intermediate risk recurrence score (RS) from a 21-gene expression assay had similar outcomes, regardless of whether they received chemotherapy, a new study from The University of Texas MD Anderson Cancer finds.

The encouraging research, published in the journal *Cancer*, still needs to be validated in an ongoing international trial. If verified, [women](#) with intermediate scores may one day be able to avoid chemotherapy as standard of care.

Historically, the management of women with early-stage [breast cancer](#) has been aggressive, explained Carlos H. Barcenas, M.D., assistant professor, Breast Medical Oncology, with many Stage I and all Stage II [patients](#) receiving chemotherapy.

Fortunately, over the last decade, research discoveries across the breast cancer landscape have evolved such that less is now more, for women with early-stage disease, said Barcenas.

"Through years of research discoveries, it became clear that we were over-treating many women with breast cancer, especially those with early-stage disease. In addition to chemotherapy's obvious side effects, there were also long-term complications for these women as survivors," said Barcenas, the study's corresponding author.

One such practice-changing clinical discovery was that a 21-gene-expression assay could predict the risk of recurrence among some with [early-stage breast cancer](#). In 2015, initial results from the international clinical trial, TAILORx, found that women with hormone receptor positive, HER2 and lymph node negative early stage disease with a low RS of 0-10 could have chemotherapy omitted altogether.

While the findings dramatically changed care for women with a low RS, many questions remain regarding the management of women with an intermediate RS, defined by this trial as a score of 11-25. The conversation to give or not give chemotherapy is one Barcenas often has in his clinic, he said.

"This study evolved out of my patients' frustration that I could not provide a definitive answer to their paramount question: 'Do I need chemotherapy?' As we wait for the TAILORx findings, we decided to look at MD Anderson's own experience to glean insight on how best to care for these women with an intermediate risk recurrence score," said Barcenas.

For the retrospective, single-institution analysis, Barcenas and his MD Anderson colleagues identified 1,424 Stage I and II [breast cancer patients](#) treated at the institution between 2005 and 2011. All underwent the 21-gene expression assay. In addition, the women were hormone receptor positive, HER2 and lymph node negative. None of the MD Anderson patients identified participated in the TAILORx trial.

Barcenas and his colleagues used the same RS cutoff scores as the TAILORx trial. Of the MD Anderson patients, the RS distribution was: 297 (21 percent) scored 0 - 10; 894 (63 percent) scored 11-25; and 233 (16 percent) scored greater than 25. Of those groups, 1.7, 15 and 73.4 percent received chemotherapy, respectively.

With a median follow-up of 58 months, those with a RS of 11-25 had an invasive disease-free survival (IDFS) rate at five years of 92.6 percent, regardless of whether patients received chemotherapy or not.

Among those patients who did not receive chemotherapy, the estimated rates of IDFS and overall survival was 93 percent and 98 percent, respectively, which was comparable to those who did receive chemotherapy.

The study has a number of limitations, said Barcnas. Due to a relatively short follow-up and the few number of outcome events, the researchers feel that the benefit of chemotherapy cannot be ruled out yet in this group of patients.

While not practice changing, the findings do give Barcnas more information when discussing the benefits of [chemotherapy](#) in early-stage breast cancer patients, should their RS score be 11-25.

"Our research is likely the most comprehensive data we will have about this patient population until the international randomized trial is published," said Barcnas. "Hopefully our findings will serve as a discussion point between physicians and patients as they are making critical decisions regarding a woman's breast cancer care."

More information: Bobby A. Tajudeen et al. Dynamic optical contrast imaging as a novel modality for rapidly distinguishing head and neck squamous cell carcinoma from surrounding normal tissue, *Cancer* (2016). [DOI: 10.1002/cncr.30338](https://doi.org/10.1002/cncr.30338)

Provided by University of Texas M. D. Anderson Cancer Center

Citation: More patients with early-stage breast cancer may be able to avoid chemotherapy in the future (2017, February 15) retrieved 8 May 2024 from <https://medicalxpress.com/news/2017-02-patients-early-stage-breast-cancer-chemotherapy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.