


Many stroke patients do not receive life-saving therapy

February 23 2017



Credit: American Heart Association

Although tPA treatment for stroke is increasing over time, minorities, women and residents of 11 southeastern states that make up the "Stroke Belt" are left behind when it comes to receiving tPA, according to research presented at the American Stroke Association's International Stroke Conference 2017.

Tissue plasminogen activator, or tPA, is the only treatment approved by the Food and Drug Administration for ischemic stroke, the most common kind of stroke. If administered within 4.5 hours of the first signs of stroke, tPA can dissolve the blood clot and restore blood flow to the affected part of the brain.

"Hospitals, governments and other organizations are undertaking efforts to increase the number of [patients](#) who receive tPA," said Tracy Madsen, M.D., Sc.M., lead researcher and Assistant Professor of Emergency Medicine at Brown University in Rhode Island. "We wanted to see if these quality improvement efforts were making a difference."

The study reviewed records from the National Inpatient Sample of 563,087 patients (median age 74) who had an [ischemic stroke](#) between 2005 and 2011. Overall, 3.8 percent of patients received tPA, with the number growing each year.

Researchers found:

- Blacks were 38 percent less likely than whites to receive tPA.
- Hispanics were 25 percent less likely than whites to receive tPA.
- Women were 6 percent less likely than men to receive tPA.
- Those with private insurance were 29 percent more likely to receive tPA compared to those with Medicare.
- Residents of the "Stroke Belt" were 31 percent less likely than those living elsewhere to receive tPA.

Researchers also found that patients discharged from a designated stroke center or a hospital participating in the American Heart Association's Get With The Guidelines—Stroke program were more likely to receive tPA. Likewise, patients were more likely to receive tPA at large, urban, or teaching hospitals compared to hospitals discharged from small, rural, or non-teaching hospitals.

Madsen said that the growing number of hospitals participating in the Get With The Guidelines—Stroke program and legislation requiring emergency services to take [stroke patients](#) to regional stroke centers are likely to increase the number of patients receiving tPA.

"Some previous studies have found that up to three-fourths of patients arrived after the time window for tPA had closed," Madsen said. "Many patients across all groups do not arrive at the hospital in time, but this is particularly true for underrepresented minorities."

An important limitation of the study data is that researchers could not determine why patients did not receive tPA. The study is also limited because we were not able to adjust for patient level factors such as time to arrival and other tPA exclusion criteria, stroke severity, patient education, and socioeconomic status. "More research needs to be done to help figure out why many patients do not receive tPA," Madsen said.

"There is also a lot of work to do in the realm of stroke education so that patients recognize stroke symptoms and call EMS immediately," Madsen said.

According to the American Heart Association's Heart Disease and Stroke Statistical Update, 795,000 Americans have a stroke every year, causing almost 129,000 deaths. Residents of the Stroke Belt—Alabama, Arkansas, Georgia, Indiana, Kentucky, Louisiana, Mississippi, North Carolina, South Carolina, Tennessee and Virginia—suffer even higher rates of stroke and [stroke](#) death.

The National Inpatient Sample is the largest publicly available database of inpatient health care in the U.S.

Provided by American Heart Association

Citation: Many stroke patients do not receive life-saving therapy (2017, February 23) retrieved 13 August 2024 from <https://medicalxpress.com/news/2017-02-patients-life-saving-therapy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.