

How does penis length change after prostate cancer surgery?

February 10 2017



Micrograph showing prostatic acinar adenocarcinoma (the most common form of prostate cancer) Credit: Wikipedia, [CC BY-SA 3.0](https://creativecommons.org/licenses/by-sa/3.0/)

Many patients who have their prostate glands removed as a treatment for prostate cancer complain of shortening in the length of the penis.

A new study conducted in 102 men found that certain anatomical changes occur that make the length shortest 10 days after surgery, but it gradually recovers to normal values after 12 months.

"The findings can help inform patients about changes in penile appearance after [radical prostatectomy](#)," said Dr. Yoshifumi Kadono, lead author of the *BJU International* study.

More information: Yoshifumi Kadono et al, Changes in penile length after radical prostatectomy: investigation of the underlying anatomical mechanism, *BJU International* (2017). DOI: [10.1111/bju.13777](https://doi.org/10.1111/bju.13777)

Provided by Wiley

Citation: How does penis length change after prostate cancer surgery? (2017, February 10) retrieved 12 May 2024 from <https://medicalxpress.com/news/2017-02-penis-length-prostate-cancer-surgery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.