

Surgery can benefit T2DM control for patients with lumbar stenosis

February 2 2017



(HealthDay)—Successful lumbar surgery is associated with improved

glycemic control for patients with lumbar spinal stenosis (LSS) and type 2 diabetes mellitus (DM-2), according to a study published in the February issue of *The Spine Journal*.

In a prospective observational study, Kyoung-Tae Kim, M.D., Ph.D., from Kyungpook National University Hospital in Daegu, South Korea, and colleagues examined the effects of postoperative improvement in physical activity on [body mass index](#) (BMI) and hemoglobin A1c (HbA1c) in patients with LSS and DM-2. One hundred nineteen patients were enrolled and assessed over a two-year follow-up period.

The researchers found that the overall HbA1c values were 7.08 ± 0.94 , 6.58 ± 0.87 , 6.59 ± 0.79 , and 6.59 ± 0.79 percent, respectively, before, and at six months, one year, and two years after surgery (P values: six months, 0.024; one year, 0.021, and two years, 0.038). The difference between pre- and postoperative HbA1c was statistically significant in the not well-controlled sugar group (P

"The present study demonstrates that in patients with DM-2 and LSS, successful lumbar surgery may facilitate [glycemic control](#) by enabling an increase in the patient's level of [physical activity](#)," the authors write.

More information: [Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Surgery can benefit T2DM control for patients with lumbar stenosis (2017, February 2) retrieved 4 May 2024 from <https://medicalxpress.com/news/2017-02-surgery-benefit-t2dm-patients-lumbar.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.