


New study shows tissue healing response following a heart attack

February 8 2017, by Matt McGowan



Heart Attack

Myocardial Infarction or Heart Attack. Credit: Blausen Medical Communications/Wikipedia/CC-A 3.0

In the weeks following a heart attack, the injured heart wall acquires more collagen fibers that are significantly less stiff due to a lack of fiber crosslinks, according to a new study by a University of Arkansas

researcher and his colleagues at Tufts University. The changes to the cardiac tissue may lead to excessive scar formation and ultimately heart failure.

The discovery was made by a team that included Kyle Quinn, assistant professor of biomedical engineering at the U of A, and Irene Georgakoudi and Lauren Black, associate professors of biomedical engineering Tufts. Their findings were published in Nature Publishing Group's *Scientific Reports*.

Using multiphoton microscopy, a powerful imaging technique, the researchers examined changes in composition and mechanical properties of the heart wall in the weeks following a [heart attack](#). Through a process of tissue decellularization, they studied the fibrous load-bearing microstructure around cardiac cells and found that following a heart attack the newly-forming scar tissue was made up of collagen fibers that were thinner, less naturally fluorescent, and more aligned than in healthy tissue. These fiber properties measured with multiphoton microscopy were associated with a poor mechanical response compared to healthy [cardiac tissue](#).

"With multiphoton microscopy, we can visualize both collagen fiber organization and crosslinking status," Quinn said. "Clearly, after a heart attack, the extra-cellular matrix was less stiff due to the deposition of new [collagen fibers](#) that lacked crosslinks."

The ability to predict the mechanical environment of the heart wall through these imaging techniques may help researchers understand how scarring, after a heart attack can ultimately lead to [heart failure](#), Quinn said. Quinn will continue working on the quantitative imaging methods to understand the relationship between the structure and mechanical function of biological tissues. His goal is to apply these methods to problems with skin and wound healing.

"Our skin shares many similar characteristics with the heart in its healing response following an ischemic injury," Quinn said.

More information: Kyle P. Quinn et al. Optical metrics of the extracellular matrix predict compositional and mechanical changes after myocardial infarction, *Scientific Reports* (2016). [DOI: 10.1038/srep35823](https://doi.org/10.1038/srep35823)

Provided by University of Arkansas

Citation: New study shows tissue healing response following a heart attack (2017, February 8) retrieved 2 May 2024 from <https://medicalxpress.com/news/2017-02-tissue-response-heart.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--