

Acupuncture improves outcomes in carpal tunnel syndrome in part by remapping the brain

March 2 2017

Though the practice of acupuncture predates current understanding of physiology by several millennia, it often provides measureable improvements in health outcomes, particularly in the area of chronic pain. Now, in a study reported in the journal *Brain*, a team of investigators based at the Athinoula A. Martinos Center for Biomedical Imaging at Massachusetts General Hospital (MGH) sheds new light on the question of how.

"Acupuncture is a medical therapy that originated in China several thousand years ago," said Vitaly Napadow, PhD, director of the Center for Integrative Pain Neuroimaging at the Martinos Center and senior author of the *Brain* paper. "But despite its long history, the intervention itself - particularly when coupled with electrical stimulation - has significant similarities to many conventional therapies, such as transcutaneous electrical nerve stimulation (TENS). A large body of clinical research exploring acupuncture for chronic pain disorders has demonstrated that it may be marginally better than a placebo procedure in reducing pain ratings. But questions still remain: How exactly does acupuncture work? Is it any better at improving objective outcomes for chronic pain?"

To get to the root of these questions, Napadow and colleagues performed a sham-controlled acupuncture neuroimaging study of carpal tunnel syndrome (CTS), a neuropathic pain disorder. Few chronic pain

disorders have established biomarkers or measureable treatment outcomes. However, in CTS measurements taken at the wrist of the speed at which signals are transmitted along the median nerve are a well known and accepted biomarker. In addition, studies by Napadow and others have shown that the [brain](#) - particularly the primary somatosensory cortex, which receives signals related to the sense of touch - is remapped in CTS. Specifically, brain cells that usually respond to touch signals from specific fingers start to respond to signals from multiple fingers, which provides another measureable outcome.

Using functional magnetic resonance imaging (fMRI) taken before and after several months of therapy in three different groups of CTS patients - one receiving electro-acupuncture at the affected hand, one receiving electro-acupuncture at the ankle opposite the affected hand, and the other receiving sham electro-acupuncture with placebo needles near the affected hand - the researchers found that both real and [sham acupuncture](#) improved patient-reported CTS symptoms. However, there were notable differences in physiologic measures. Real acupuncture at the affected hand led to measurable improvements in outcomes both at the affected wrist and in the brain, while acupuncture at the opposite ankle produced improvement at the wrist only. Brain remapping immediately after real acupuncture was linked to long-term improvement in CTS symptoms. No physiologic improvements resulted from sham acupuncture.

Even after years of clinical research, controversy continues as to whether acupuncture works primarily by the placebo effect, especially given the slight differences between the efficacy of real and sham acupuncture. The findings of the *Brain* study help to address this question. Sham acupuncture may produce a stronger placebo effect than a pill because it sends inputs to the brain via skin receptors and is coupled with a specific ritual. But the symptom improvement produced by sham treatment for conditions like CTS also might derive from entirely different

mechanisms than those elicited by real acupuncture, the mechanisms of which may more specifically target CTS pathophysiology.

"Sham acupuncture may 'work' by modulating known placebo circuitry in the brain," Napadow said. "In contrast, real acupuncture may improve CTS symptoms by rewiring the primary somatosensory cortex, in addition to modulating local blood flow to the peripheral nerve in the wrist. In other words, both peripheral and central neurophysiological changes in CTS may be halted or even reversed by electro-acupuncture interventions that provide more prolonged and regulated input to the brain - something that future, longer-term neuroimaging studies should explore."

Napadow and colleagues plan to follow up the *Brain* study with further research linking objective/physiological and subjective/psychological outcomes for acupuncture-produced pain relief. Better understanding of how [acupuncture](#) works to relieve pain ultimately will enable them and others to optimize the therapy to provide effective, non-pharmacological care for [chronic pain](#) patients.

More information: Yumi Maeda et al, Rewiring the primary somatosensory cortex in carpal tunnel syndrome with acupuncture, *Brain* (2017). [DOI: 10.1093/brain/awx015](https://doi.org/10.1093/brain/awx015)

Provided by Massachusetts General Hospital

Citation: Acupuncture improves outcomes in carpal tunnel syndrome in part by remapping the brain (2017, March 2) retrieved 27 April 2024 from <https://medicalxpress.com/news/2017-03-acupuncture-outcomes-carpal-tunnel-syndrome.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.