

Antidepressants and anti-anxiety drugs linked to increased risk of hip fracture

March 8 2017



Credit: Susan Buck Ms/Public Domain

In a recent study, older individuals who had fallen and broken a hip used antidepressant and anti-anxiety medications more frequently than the

general older population.

The analysis measured [drug](#) levels in individuals' [blood samples](#), which provided investigators with a more precise measure than asking participants if they took the medications or examining their medical records. Indeed, the drugs were often present in these samples despite no information in patients' [medical records](#).

"We were surprised to find that so many patients had centrally-acting drugs in their blood without doctors being aware of this use. Several of the detected drugs increase the risk of falling, which underscores the clinical relevance of these findings," said Ragnhild Birkeland Waade, lead author of the *British Journal of Clinical Pharmacology* study.

More information: Ragnhild Birkeland Waade et al, Psychotropics and weak opioid analgesics in plasma samples of older hip fracture patients - detection frequencies and consistency with drug records, *British Journal of Clinical Pharmacology* (2017). [DOI: 10.1111/bcp.13244](#)

Provided by Wiley

Citation: Antidepressants and anti-anxiety drugs linked to increased risk of hip fracture (2017, March 8) retrieved 2 May 2024 from <https://medicalxpress.com/news/2017-03-antidepressants-anti-anxiety-drugs-linked-hip.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
