


Some blood thinners may increase heart attack risk

March 22 2017



Heart Attack

Myocardial Infarction or Heart Attack. Credit: Blausen Medical Communications/Wikipedia/CC-A 3.0

A new study has examined whether different blood thinning medications prescribed to prevent strokes in patients with atrial fibrillation might increase the risk of heart attacks.

In the retrospective study of 30,146 patients, investigators found a twofold increased risk in patients taking direct acting oral anticoagulants (dabigatran and rivaroxaban) compared with those taking vitamin k antagonists such as warfarin. Heart attack risk was also higher in patients taking aspirin than in those taking warfarin.

"More research should be on-going as use of direct acting oral anticoagulants increases in the population," wrote the authors of the *British Journal of Clinical Pharmacology* study.

More information: *British Journal of Clinical Pharmacology*, [DOI: 10.1111/bcp.13264](https://doi.org/10.1111/bcp.13264)

Provided by Wiley

Citation: Some blood thinners may increase heart attack risk (2017, March 22) retrieved 26 April 2024 from <https://medicalxpress.com/news/2017-03-blood-thinners-heart.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--