

# Childhood adiposity linked to later risk of fatty liver disease

March 29 2017

---



(HealthDay)—Childhood adiposity is associated with increased odds of

nonalcoholic fatty liver disease (NAFLD), according to a study published online March 29 in *Pediatrics*.

Yinkun Yan, Ph.D., from the Capital Institute of Pediatrics in Beijing, and colleagues followed 1,350 subjects (aged 28 to 45 years) from 2010 to 2014 from an original cohort of children aged 6 to 18 years. Body mass index (BMI) and subscapular skinfold thickness were used to define childhood overweight and obesity.

The researchers found that, compared with normal-weight children, overweight and [obese children](#) were more likely to have adult NAFLD (males: odds ratios, 2.49 and 2.78 for BMI and subscapular skinfold thickness, respectively; females: odds ratios, 3.34 and 3.61, respectively; all P alanine aminotransferase (ALT) (males: odds ratios, 1.64 and 1.66, respectively; females: odds ratios, 2.12 and 3.01, respectively; all P childhood obesity, compared with those who had normal weight in childhood and were nonobese in adulthood. The likelihood of having NAFLD and ALT elevation in adulthood for both sexes was similar for subjects who were overweight or obese in childhood but became nonobese in adulthood.

"The risk associated with increased weight during childhood can be mitigated by becoming nonobese in [adulthood](#)," the authors write.

**More information:** [Abstract/Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Childhood adiposity linked to later risk of fatty liver disease (2017, March 29) retrieved 30 April 2024 from <https://medicalxpress.com/news/2017-03-childhood-adiposity-linked-fatty-liver.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.