


Early intervention with new treatment enables durable control of HIV-like virus in monkeys

March 13 2017



This protein-structure model shows how the broadly neutralizing antibodies 10-1074 (blue) and 3BNC117 (red) that were infused into the monkeys each

bind to a different site on the SHIV spike (beige and pink), helping prevent the virus from escaping the antibodies. Credit: Gristick et al., *Nature Structural & Molecular Biology* 2016.

There are more than 25 drugs to control HIV, yet the virus remains one of the world's biggest health problems. One of the many challenges with existing therapies is that a dormant version of the virus is always lurking in the background, ready to attack the immune system as soon as treatment is interrupted.

Now, new research from The Rockefeller University and the National Institutes of Health suggests that treatment with two anti-HIV antibodies immediately after infection enables the [immune system](#) to effectively control the virus, preventing its return for an extended period.

"This form of therapy can induce potent immunity to HIV, allowing the host to control the infection," says Michel Nussenzweig, head of the Laboratory of Molecular Immunology and an Investigator with the Howard Hughes Medical Institute. "It works by taking advantage of the immune system's natural defenses, similar to what happens in some forms of cancer immunotherapy."

The research was conducted in [macaque monkeys](#), using a model of HIV infection called [simian-human immunodeficiency virus](#) (SHIV).

Although this model does not precisely mimic human HIV infection, the findings suggest that immunotherapy should be explored as a way of controlling the virus and boosting an [immune response](#) that might be capable of controlling the infection in people. The study publishes on March 13 in *Nature*.

Long-term control

The two drugs used in the study, [3BNC117](#) and [10-1074](#), belong to a class of molecules called broadly neutralizing antibodies. They were discovered by the Nussenzweig laboratory in studies of "elite controllers," people whose immune systems have a rare ability to fight off the virus. Each antibody binds to a different site of the virus, preventing its damaging effects from different angles.

13 monkeys were inoculated with the SHIV virus, and then given three intravenous infusions of the two antibodies over a two-week period. The treatment suppressed the virus to levels near or below the limit of detection, and its effect lasted for as long as six months. After the antibodies had cleared out of the monkeys' bodies, the virus rebounded in all but one animal.

But then, 5 to 22 months later, something remarkable happened: six of the monkeys spontaneously regained control of the virus. Their virus levels once again plummeted to undetectable levels and remained suppressed for another 5 to 13 months.

These six monkeys were also able to maintain healthy levels of key immune cells after receiving the antibody infusions.

In addition, four other monkeys that did not regain complete control of the virus nevertheless showed promising responses to the treatment: they maintained extremely low viral loads and healthy levels of key immune cells for two to three years after infection. In total, 10 of the 13 monkeys benefitted from antibody immunotherapy.

Feasibility in humans

Nussenzweig and colleagues also investigated what aspect of the immune system was helping the monkeys ward off the virus's return. They gave

the six controller monkeys an antibody that targets and depletes a type of [immune](#) cell called cytotoxic T cells. Infusion of this antibody immediately increased the amount of SHIV in the [monkeys'](#) blood and decreased cytotoxic T cell levels, indicating that these cells play a key role in preventing SHIV replication after therapeutic antibody infusion.

The researchers are now repeating this experiment after a longer exposure to the [virus](#), waiting two to six weeks after SHIV infection before administering the therapeutic antibody infusions. This is how long it usually takes for an HIV-infected person to be diagnosed and able to receive treatment.

More information: Early antibody therapy can induce long-lasting immunity to SHIV, [nature.com/articles/doi:10.1038/nature21435](https://doi.org/10.1038/nature21435)

Provided by Rockefeller University

Citation: Early intervention with new treatment enables durable control of HIV-like virus in monkeys (2017, March 13) retrieved 28 April 2024 from <https://medicalxpress.com/news/2017-03-early-intervention-treatment-enables-durable.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|