

Endocrine Society experts issue Clinical Practice Guideline on hypothalamic amenorrhea

March 22 2017

Female athletes and women who have eating disorders are prone to developing a condition called hypothalamic amenorrhea that causes them to stop menstruating. The Endocrine Society today issued a Clinical Practice Guideline advising healthcare providers on ways to diagnose and treat this condition.

The guideline, titled "Functional Hypothalamic Amenorrhea: An Endocrine Society Clinical Practice Guideline," was published online and will appear in the May 2017 print issue of *The Journal of Clinical Endocrinology & Metabolism (JCEM)*, a publication of the Endocrine Society.

Hypothalamic amenorrhea occurs when the hypothalamus in the brain slows or stops releasing GnRH, the hormone that controls the [menstrual cycle](#). It often affects adolescent girls or women with low body weight, a low percentage of body fat, a very low calorie or fat intake, and emotional stress. Ballet dancers, figure skaters, runners and others who burn more calories through exercise than they consume in their diet can be at risk for developing hypothalamic amenorrhea.

"This energy imbalance needs to be addressed to effectively treat hypothalamic amenorrhea and typically requires behavioral modifications," said Catherine M. Gordon of Cincinnati Children's Hospital Medical Center in Cincinnati, Ohio, and the chair of the task

force that authored the guideline. "Referring patients to a nutritionist for specialized dietary instructions is an extremely important part of their care. Menstrual cycles can often be restored with increased calorie consumption, improved nutrition or decreased exercise activity."

Hypothalamic amenorrhea raises the risk of other health problems, such as delayed puberty in adolescents and infertility in adult women. Chronic hypothalamic amenorrhea can contribute to bone loss, and some patients develop stress fractures and are at high risk to develop osteoporosis.

To diagnose hypothalamic amenorrhea, [healthcare providers](#) must rule out other conditions that can halt menstruation, including benign tumors in the [pituitary gland](#) and adrenal gland disorders. The guideline recommends that providers first exclude pregnancy as a cause and then perform a full physical exam to evaluate for other potential causes. General laboratory tests, including a complete blood count and electrolytes, also are part of the recommended screening process.

Recommendations from the guideline include:

- Hypothalamic amenorrhea is a "diagnosis of exclusion," which requires healthcare providers to rule out other conditions that could be interrupting the menstrual cycle.
- As part of their initial evaluation, women diagnosed with hypothalamic amenorrhea should have a series of laboratory tests to check levels of hormones including estrogen, thyroid hormones and prolactin. The workup can help identify factors preventing menstruation.
- Hypothalamic amenorrhea patients should be evaluated for inpatient treatment if they have an abnormally slow heart rate, low blood pressure, or an electrolyte imbalance. Careful monitoring is needed in these cases because there is a high mortality rate associated with hypothalamic amenorrhea in the

setting of eating disorders, particularly anorexia nervosa.

- Select patients presumed to have hypothalamic [amenorrhea](#) should undergo a brain MRI to check for damage to or abnormalities of the pituitary gland or pituitary hormone deficiencies, if they exhibit select signs or symptoms, including a history of severe or persistent headaches; persistent vomiting that is not self-induced; changes in vision, thirst or urination not attributable to other causes; neurological signs suggesting a central nervous system abnormality; or other clinical signs or test results that suggest pituitary hormone deficiency or excess.

More information: Hypothalamic Amenorrhea: An Endocrine Society Clinical Practice Guideline, [DOI: 10.1210/jc.2017-0131](#) , [academic.oup.com/jcem/article-... irectedFrom=fulltext](#)

Provided by The Endocrine Society

Citation: Endocrine Society experts issue Clinical Practice Guideline on hypothalamic amenorrhea (2017, March 22) retrieved 2 May 2024 from <https://medicalxpress.com/news/2017-03-endocrine-society-experts-issue-clinical.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--