

Repeated eye injections for age-related macular degeneration associated with increased risk for glaucoma

March 16 2017

Patients with age-related macular degeneration who received seven or more eye injections of the drug bevacizumab annually had a higher risk of having glaucoma surgery, according to a study published online by *JAMA Ophthalmology*.

The advent of intravitreal (in the vitreous, the fluid behind the lens in the eye) anti-vascular endothelial growth factor (VEGF) injections to treat common causes of vision loss, such as exudative (wet) age-related macular degeneration (AMD) and [diabetic macular edema](#) has improved visual outcomes for many patients. Intravitreal injections of anti-VEGF agents may increase the risk of elevated intraocular pressure (IOP); however, the risk of developing moderate to advanced glaucoma requiring [glaucoma surgery](#) has been unclear.

Brennan D. Eadie, M.D., Ph.D., of the University of British Columbia, Vancouver, and colleagues conducted a study that included patients who had received intravitreal bevacizumab injections for exudative [age-related macular degeneration](#). Cases were identified using glaucoma surgical codes for various procedures. For each case, 10 controls were identified; the number of intravitreal bevacizumab injections received per year was determined for both cases and controls.

Seventy-four cases of glaucoma surgery and 740 controls were identified. The researchers found that seven or more injections were

associated with a significantly higher risk of glaucoma surgery.

"Clinicians should be aware of the potential association of repeated, recent intravitreal anti-VEGF injections for diseases, such as exudative AMD, with subsequent need for glaucoma surgery," the authors write.

More information: *JAMA Ophthalmol.* Published online March 16, 2017. [DOI: 10.1001/jamaophthalmol.2017.0059](https://doi.org/10.1001/jamaophthalmol.2017.0059)

Provided by The JAMA Network Journals

Citation: Repeated eye injections for age-related macular degeneration associated with increased risk for glaucoma (2017, March 16) retrieved 25 April 2024 from <https://medicalxpress.com/news/2017-03-eye-age-related-macular-degeneration-glaucoma.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|