

Musculoskeletal symptoms predict psoriatic arthritis

March 13 2017



(HealthDay)—For patients with psoriasis, nonspecific musculoskeletal



symptoms, including joint pain, fatigue, and stiffness, predict the development of psoriatic arthritis (PsA), according to a study published in the March issue of *Arthritis & Rheumatology*.

Lihi Eder, M.D., Ph.D., from the University of Toronto, and colleagues conducted a <u>prospective cohort study</u> involving patients with psoriasis who were assessed at baseline and reassessed annually to examine development of clinical PsA. At each visit, the presence of musculoskeletal symptoms and the patients' assessments of pain, fatigue, stiffness, physical function, and psychological distress were recorded.

The researchers found that 57 of 410 patients with psoriasis developed PsA. Prediction of subsequent PsA was associated with baseline presence of arthralgia in women (hazard ratio [HR], 2.59), heel pain (HR, 4.18), high fatigue score (HR, 2.36), and high stiffness score (HR, 2.03). Increases from baseline in fatigue, pain, and stiffness scores (HRs, 1.27, 1.34, and 1.21, respectively), and a worsening in physical function score (HR, 0.96) also predicted PsA development.

"A preclinical phase exists in patients with PsA prior to the diagnosis of the disease," the authors write. "This phase is characterized by nonspecific musculoskeletal symptoms, including joint pain, fatigue, and stiffness."

More information: Abstract

Full Text (subscription or payment may be required)

Copyright © 2017 HealthDay. All rights reserved.

Citation: Musculoskeletal symptoms predict psoriatic arthritis (2017, March 13) retrieved 19 April 2024 from

https://medicalxpress.com/news/2017-03-musculoskeletal-symptoms-psoriatic-arthritis.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.