

## Proton therapy offers new treatment possibility for recurrent lung cancer

## March 16 2017

A new study offers hope for patients with recurrent lung cancer, who historically have been considered ineligible for curative treatment. In the largest analysis to date of reirradiation using intensity-modulated proton therapy (IMPT) for lung and other thoracic tumors, more than three-fourths of patients were free from local recurrence at one year following retreatment, and fewer than one in ten patients experienced severe side effects. The study will be presented today at the <u>2017 Multidisciplinary</u> <u>Thoracic Cancers Symposium</u>.

Lung cancer causes more deaths in the United States than any other type of cancer, due in part to its aggressive nature and likelihood of recurrence. Historically, recurrences have been challenging to treat, because many of these patients are not candidates for surgery. Moreover, concerns about cumulative <u>radiation</u> doses to essential organs near thoracic tumors (e.g., heart, lungs, esophagus) may limit the use of curative radiation therapy (RT) for patients who received thoracic RT in the past. Accordingly, patients whose lung cancer recurs are generally offered only palliative options to manage pain and other symptoms.

Proton therapy, an advanced type of RT, allows radiation oncologists to spare critical normal tissues while delivering escalated, curative doses of radiation to nearby tumors. In particular, IMPT is able to exactly target a tumor, even if the tumor wraps around a critical normal tissue structure, such as a lung.

"Treating patients who have already received a prior course of thoracic



radiation is a common clinical scenario, and it is particularly challenging to subsequently provide strong enough radiation doses to eliminate the new tumor without causing significant harm to normal tissues," said Jennifer Ho, MD, lead author of the study and a resident in <u>radiation</u> <u>oncology</u> at the University of Texas MD Anderson Cancer Center in Houston. "Our study is the first to show that IMPT can be safe and effective for these patients—and that it offers these patients a chance for lasting cancer control without adding significant toxicity."

Researchers retrospectively examined the records of 27 patients who received reirradiation for thoracic tumors using the IMPT technique through prospective clinical trials at a single institution between 2011 and 2016. Twenty-two patients (81%) had non-small cell <u>lung cancer</u> (NSCLC). All patients had previously received curative thoracic RT.

Time to reirradiation ranged from 0.1 months to 212 months, with a median of 29.5 months. The median radiation dose was 66 Gray (Gy), with a range of 43.2 to 84 Gy, and it was delivered in 2 Gy fractions. The median follow-up for all patients in the study was 11.2 months.

The median overall survival (OS) for patients in this study was 18 months following IMPT reirradiation. At one year following retreatment, the majority of patients were free from local and regional relapse (freedom from local failure = 78%; local-regional relapse = 61%). Just over half of the patients were free from disease progression (progressionfree survival, PFS = 51%), and the one-year OS rate was 54 percent. Four of the patients (15%) experienced a local recurrence within one year of retreatment.

Patients who were prescribed a higher dose of IMPT reirradiation were even less likely to experience recurrence or progression. At one-year follow-up, patients who received IMPT reirradiation doses at or above the population median were twice as likely to be free from local failure



(100% vs. 49%, p = 0.01) and nearly four times as likely to be free from local-regional failure (84% vs. 23%, p = 0.035). The one-year PFS rate was five times higher for these patients, as well (76% vs. 14%, p = 0.05). Several disease characteristics, namely higher T stage at diagnosis, squamous histology and higher recurrent tumor volume, were associated with worse OS rates.

Reirradiation with IMPT was well-tolerated among the patients. Only two patients (7%) experienced moderate to severe long-term lung toxicity (i.e., grade 3 or higher <u>side effects</u>). There were no severe longterm esophageal side effects nor any life-threatening toxicities among these patients.

"Historically, 20 to 30 percent of <u>patients</u> have experienced moderate or severe side effects, and even fatal side effects, following reirradiation. We knew that IMPT would allow us to generate much more precise radiation treatment plans that spared normal tissue, but we weren't sure if this would translate into excellent clinical outcomes," said Joe Y. Chang, MD, senior author of the study and a professor of radiation oncology at MD Anderson. "Our findings demonstrate that the use of IMPT resulted in better local control and survival, with very minimal toxicity, compared to other radiation types, suggesting that IMPT is the optimal treatment modality for re-treatment of thoracic cancers."

## Provided by American Society for Radiation Oncology

Citation: Proton therapy offers new treatment possibility for recurrent lung cancer (2017, March 16) retrieved 6 May 2024 from <u>https://medicalxpress.com/news/2017-03-proton-therapy-treatment-possibility-recurrent.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.