

Study shows role of depression in the ongoing tobacco epidemic

March 1 2017

The prevalence of smoking has remained fairly stable over the past decade after declining sharply for many years. To determine whether an increase in certain barriers to successful cessation and sustained abstinence may be contributing to this slowed decline, researchers at Columbia University's Mailman School of Public Health analyzed changes in the prevalence of depression among current, former and never smokers in the U.S. The team found that depression appeared to have significantly increased in the U.S. from 2005 to 2013 among smokers, as well as among former and never smokers. While the prevalence of depression is consistently highest among smokers, the rate of increase in depression was most prominent among former and never smokers. The full study findings are published online in the journal *Drug and Alcohol Dependence*.

The research team, led by Renee Goodwin, PhD, in the Department of Epidemiology, analyzed data from the National Household Survey on Drug Use, an annual cross-sectional study of approximately 497,000 Americans, ages 12 and over. The prevalence of past 12-month depression was examined annually among current (past 12-month), former (not past 12-month), and lifetime non-[smokers](#) from 2005 to 2013. The researchers further analyzed the data by age, gender, and household income.

"The prevalence of depression prevalence increased and remains higher among current smokers overall, but the rate of the increase among former and never smokers was even more prominent," noted Dr.

Goodwin. Striking temporal changes emerged by age, gender and income. Specifically, depression increased significantly, from 16 percent to 22 percent, among current smokers aged 12 to 17, and the prevalence was consistently more than twice as high as that of never smokers. The increase in depression also rose from 6 percent to 8 percent among [male smokers](#) and increased from 6 percent to 9 percent among smokers in the highest income group. Throughout this period, the prevalence of depression among current smokers was consistently twice as high as among former and never smokers. "The very high rates of depression among the youngest smokers, those aged 12-17, is very concerning, as it may impair their ability not only to stop smoking, but also to navigate the important developmental tasks of adolescence that are important for a successful adult life" said Mailman School of Public Health's Dr. Deborah Hasin, a senior member of the research team.

"Public health efforts aimed at decreasing the prevalence of smoking must take depression into account, a common and modifiable barrier whose treatment may help to increase successful smoking cessation," said Dr. Goodwin, adjunct associate professor of Epidemiology. "We also need to examine factors that may be leading increases in [depression](#) in the U.S. population among both smokers and non-smokers."

Provided by Columbia University's Mailman School of Public Health

Citation: Study shows role of depression in the ongoing tobacco epidemic (2017, March 1) retrieved 25 April 2024 from <https://medicalxpress.com/news/2017-03-role-depression-ongoing-tobacco-epidemic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.