


Scientists zoom in on AIDS virus hideout

March 15 2017



HIV-1 Virus. Credit: J Roberto Trujillo/Wikipedia

French scientists said Wednesday they had found a way to pinpoint elusive white blood cells which provide a hideout for the AIDS virus in people taking anti-HIV drugs.

Being able to spot, and one day neutralise, these "reservoir" cells has long been a holy grail in the quest to wipe out AIDS and the [human immunodeficiency virus](#) (HIV) that causes it.

The discovery "paves the way to a better fundamental understanding of viral reservoirs," said France's CNRS research institute, which took part in the study published in *Nature*.

"In the longer term, it should lead to therapeutic strategies aiming to eliminate the latent virus," it added in a statement.

There is no cure for HIV, and infected people have to take virus-suppressing drugs for life.

This is because a small number of [immune system cells](#), in a category of cell called CD4 T lymphocytes, provide a haven for the virus, enabling it to re-emerge and spread if treatment is stopped— even after decades.

In tests using the blood of HIV patients, the researchers managed to spot a protein, dubbed CD32a, on the surface of virus-infected reservoir cells.

It was absent from [healthy cells](#).

Such a "marker" has proved very difficult to find, explained AIDS researcher Douglas Richman from the University of California San Diego, who did not take part in the study.

A person infected with HIV has about 200 billion CD4 T cells, of which

only one in a million act as virus reservoirs.

Two percent of the body's CD4 T cells (some four billion) are found in the approximately five litres of blood in an adult human, said Richman.

This means that a 100-millilitre blood sample would contain about 80 million CD4 T cells, of which around 80 would be virus reservoirs.

Whether CD32a plays an active part in enabling the [virus](#) to hole up in CD4 cells is a big question.

If so, it could throw open a tempting target for drugs to block the stealthy process.

While describing the study as "potentially seminal", Richman cautioned that CD32a was a marker found in only about half of CD4 T reservoir cells.

To eradicate latent HIV would require the targeting of a much larger proportion.

It also remains to be seen whether CD32a is as good a marker for non-blood CD4 T [cells](#) in the lymph nodes, bone marrow, gut and other tissues which could be reservoirs, he added.

More information: Benjamin Descours et al. CD32a is a marker of a CD4 T-cell HIV reservoir harbouring replication-competent proviruses, *Nature* (2017). [DOI: 10.1038/nature21710](https://doi.org/10.1038/nature21710)

© 2017 AFP

Citation: Scientists zoom in on AIDS virus hideout (2017, March 15) retrieved 1 May 2024 from

<https://medicalxpress.com/news/2017-03-scientists-aids-virus-hideout.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.