

Supervised self-injection ups teens' comfort with approach

March 16 2017



(HealthDay)—For food-allergic adolescents at risk for anaphylaxis,

supervised self-injection with an empty syringe is associated with improved comfort levels with self-injection, according to a study published in the March-April issue of the *Journal of Allergy and Clinical Immunology: In Practice*.

Eyal Shemesh, M.D., from the Icahn School of Medicine at Mount Sinai, N.Y., and colleagues randomized 60 adolescent/parent pairs to self-injection, involving supervised self-injection with an empty syringe, versus control (education only) to examine the impact on comfort and anxiety during and after clinic visits in a food allergy center.

The researchers found that supervised self-injection correlated with a significant immediate increase in the primary outcome of comfort levels with the injection (within-group comparison: mean scores, 6.93 preintervention versus 8.37 postintervention; *P* anxiety, although these differences were not statistically significant).

"A self-injection (with an empty syringe) procedure in a clinic setting improves adolescents' and parents' comfort level with self-injecting," the authors write. "It may translate into substantial clinical benefits should self-injection be needed."

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Supervised self-injection ups teens' comfort with approach (2017, March 16) retrieved 3 May 2024 from
<https://medicalxpress.com/news/2017-03-self-injection-ups-teens-comfort-approach.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.