

Alterations in soluble mediators predict transition to SLE

March 8 2017



(HealthDay)—For unaffected relatives of systemic lupus erythematosus

(SLE) patients, preclinical alterations in levels of soluble mediators may predict transition to disease, according to a study published in the March issue of *Arthritis & Rheumatology*.

Melissa E. Munroe, M.D., Ph.D., from the Oklahoma Medical Research Foundation in Oklahoma City, and colleagues examined factors associated with transition to classified [disease](#) in 409 previously identified blood [relatives](#) of patients with SLE who had up to four American College of Rheumatology (ACR) classification criteria for SLE at baseline.

The researchers found that at the time of follow-up (mean, 6.4 years), 11 percent of unaffected relatives had transitioned to classified SLE. More lupus-associated autoantibody specificities and higher SLE-specific portion of the Connective Tissue Disease Screening Questionnaire (SLE-CSQ) scores at baseline (P

"Immune perturbations precede SLE classification and can help identify high-risk relatives for rheumatology referral and potential enrollment in prevention trials," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Alterations in soluble mediators predict transition to SLE (2017, March 8) retrieved 23 April 2024 from <https://medicalxpress.com/news/2017-03-soluble-transition-sle.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
