

Researchers connect common fats to a lazy lifestyle and diabetes

April 12 2017



UBC researcher Sanjoy Ghosh connects common fats to a lazy lifestyle and diabetes. Credit: UBC Okanagan

A UBC researcher is suggesting the types of cooking oils people consume may be sabotaging their efforts to stay healthy and avoid

illnesses such as diabetes.

Sanjoy Ghosh, a Michael Smith Health Research Foundation Scholar and a professor at UBC's Okanagan campus, has recently published research that concludes a high consumption of polyunsaturated [fatty acids](#) (PUFA) but not [monounsaturated fatty acids](#) (MUFA) can lead to sedentary, in fact, lazy behaviour especially in women .

Ghosh says not that long ago, heart disease was linked to saturated fats—an idea that has become increasingly controversial in recent years. This thinking instigated the intentional removal of saturated fatty acids from most food supplies in favour of MUFA and PUFA. Essentially all fats in our 'convenience' foods like potato chips, energy bars, crackers or burgers use cooking oils like corn, sunflower and soybean and margarine—all rich in MUFAs and PUFAs.

For his research, Ghosh collaborated with UBC biologist and data analyst Jason Pither, the first-author of the study, to examine data from 21 countries in Europe. They worked specifically with data relating to pre-teen girls and then, in a second study, the [blood glucose levels](#) of [adult women](#). In putting details such as the amount of time each week spent watching TV along with other filters like a country's per capital GDP, urbanization, and even latitude, they came out with a clear connection to the consumption of n-6 polyunsaturated fatty acids and an increase in sedentary behaviour.

In particular, a significant correlation was observed in sedentary behaviour of the 11-year-old girls and PUFA in their diets.

"Our study does present new evidence that dietary PUFA is strongly associated with [sedentary behavior](#) among pre-teen girls and weakly associated with diabetes among adult women across Europe," says Ghosh, recommending for more trials and studies to confirm his

findings.

Such clinical findings come on the heels of a similar 2015 study from Ghosh's lab published in *Journal of Nutritional Biochemistry*, which provided the first indication that the omega-6 PUFAs we eat can make mice lazy.

"This data is extremely significant," says Ghosh. "Nobody has made this connection and it's time for an intervention. And if someone is beginning an exercise program without taking a close look at the fats, especially PUFA they are consuming, or changing what they're eating, then it might be doomed to failure."

More information: Jason Pither et al, Analysis using national databases reveals a positive association between dietary polyunsaturated fatty acids with TV watching and diabetes in European females, *PLOS ONE* (2017). [DOI: 10.1371/journal.pone.0173084](https://doi.org/10.1371/journal.pone.0173084)

Chi Kin Wong et al. A high-fat diet rich in corn oil reduces spontaneous locomotor activity and induces insulin resistance in mice, *The Journal of Nutritional Biochemistry* (2015). [DOI: 10.1016/j.jnutbio.2014.11.004](https://doi.org/10.1016/j.jnutbio.2014.11.004)

Provided by University of British Columbia

Citation: Researchers connect common fats to a lazy lifestyle and diabetes (2017, April 12) retrieved 4 May 2024 from <https://medicalxpress.com/news/2017-04-common-fats-lazy-lifestyle-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
