

Depressed veterans with heart disease face financial barriers to care

April 3 2017

Veterans with heart disease who are also depressed are more likely than those without depression to have trouble paying for medications and medical visits and often report delays in seeking medical care, according to research presented at the American Heart Association's Quality of Care and Outcomes Research 2017 Scientific Sessions.

Data from the Centers for Disease Control and Prevention's 2013 Behavioral Risk Factor Surveillance System survey was used to study 13,126 veterans who reported being told that they had a heart attack, stroke, or angina or coronary [heart disease](#) by a health professional.

Of those studied, 22 percent reported having been diagnosed with depression. Veteran heart patients with depression were more likely to be older and non-white, and were less likely to be employed, own a home, and had lower annual income.

They had higher rate of high blood pressure, elevated cholesterol, chronic [obstructive pulmonary disease](#) (a lung disease that causes breathing difficulty), [chronic kidney disease](#), asthma, smoking, obesity and lack of physical inactivity.

Compared to veteran heart patients without depression, those with depression were about:

- twice as likely to report difficulty affording medical care
- twice as likely to report delays in seeking medical care; and

- 45 percent more likely to report difficulty affording prescription drugs.

Data about the veteran participants in this study came from the participants' own responses, rather than from medical records. The survey did not discern whether the veterans received care at Department of Veterans Affairs (VA) facilities or other non-VA facilities.

"All [health care providers](#) who treat veterans with depression should routinely ask their patients about any difficulty with being able to pay for [medical care](#) or medications," said the study's senior author, Puja Parikh, M.D., M.P.H., an interventional cardiologist and assistant professor at Stony Brook School of Medicine and Director of Invasive Cardiology at the Northport Veterans Affairs Medical Center. "In some cases, less expensive medications can be considered and 90-day supply of drugs can be prescribed instead of a 30-day supply which can reduce costs. We can also recommend working with a social worker to obtain vouchers or coupons from pharmaceutical companies to get further financial assistance with medication costs."

Healthcare providers also should screen veterans for depression and monitor them for electrocardiogram abnormalities if they are taking anti-depressant medications, because certain psychiatric drugs can predispose some patients to develop arrhythmias. They should also counsel patients on how to take their cardiac medications, she said. Because depressed veteran heart patients are likely to have multiple chronic medical illnesses, they should be followed by a primary care doctor, psychiatrist, and cardiologist.

"Further, VA-based research could shed more light on how mental illness affects heart disease care for veterans," Parikh said. Also, future national surveys could include other mental health diagnoses that impact veterans, such as post-traumatic stress disorder (PTSD), she said.

Provided by American Heart Association

Citation: Depressed veterans with heart disease face financial barriers to care (2017, April 3)
retrieved 5 May 2024 from

<https://medicalxpress.com/news/2017-04-depressed-veterans-heart-disease-financial.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.