

Elevated biomarker following surgery linked to increased risk of death

April 25 2017

Among patients undergoing noncardiac surgery, peak postoperative high-sensitivity troponin T measurements (proteins that are released when the heart muscle has been damaged) during the first three days after surgery were associated with an increased risk of death at 30 days, according to a study published by *JAMA*.

Large observational studies suggest that among [patients](#) 45 years or older undergoing major noncardiac [surgery](#), more than 1 percent die in hospital or within 30 days of surgery. Little is known about the relationship between perioperative high-sensitivity troponin T (hsTnT) measurements and 30-day mortality and myocardial (the muscular tissue of the heart) injury after noncardiac surgery (MINS). P.J. Devereaux, M.D., Ph.D., of McMaster University, Hamilton, Ontario, Canada, and colleagues conducted a study that included patients 45 years or older who underwent inpatient [noncardiac surgery](#) and had postoperative hsTnT measurements six to 12 hours after surgery and daily for three days. The patients were recruited at 23 centers in 13 countries.

Among 21,842 participants, the average age was 63 years and 49 percent were female. Death within 30 days after surgery occurred in 266 patients. Analysis indicated that compared with the reference group (peak hsTnT

Citation: Elevated biomarker following surgery linked to increased risk of death (2017, April 25) retrieved 20 April 2024 from

<https://medicalxpress.com/news/2017-04-elevated-biomarker-surgery-linked-death.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.