

Giardiasis may be a disease of the ecology of the GI tract

April 20 2017

Colonization by the human and animal parasite, *Giardia*, changed the species composition of the mouse microbiome in a way that might be harmful. The research is published in *Infection and Immunity*, a journal of the American Society for Microbiology.

"This shift is generally characterized by more aerobic bacteria and less diversity of anaerobic species in the gastrointestinal tract," said coauthor Scott C. Dawson, Associate Professor of Microbiology and Molecular Genetics, the University of California, Davis. "This suggests that *Giardia* [infection](#) could—at least in part—be an ecological disease, with parasite colonization disrupting the previously stable ecology of the gut."

Indeed, "We found changes in the host microbial community throughout the entire gastrointestinal tract," said Dawson. These findings challenge the conventional wisdom that *Giardia* is a disease of the small intestine.

"We infected mice with *Giardia*, sacrificed them at different times post-infection, and sequenced specific regions throughout the entire gastrointestinal tract using high-throughput 16S ribosomal RNA sequencing," said Dawson. This enabled the investigators to quantify the shifts in microbial diversity in each part of the GI tract, following infection. They also pre-treated one cohort of mice with antibiotics, to determine if that pretreatment resulted in different shifts in the microbiome as compared to those in mice not receiving antibiotics (it did so).

The various investigators had different motives for their interest in conducting this research, said Dawson. Steven M. Singer, of Georgetown University, Washington, DC, had previously noted that when mice were infected with a human *Giardia*, they had to be pre-treated with antibiotics in order for a robust infection to develop. "That suggested that the gut microbiome had a protective effect in limiting giardiasis," said Dawson.

Furthermore, giardiasis is a major cause of sickness and malnutrition in developing nations, said Dawson, infecting an estimated 200 million in Africa, Asia, and Latin America, "often infecting children in communities without access to clean water. In the US, more than 15,000 cases were reported in 2012.

"While symptoms can be severe or even debilitating, the mechanism by which *Giardia* infection causes the disease symptoms remains unclear," said Dawson. "As an anaerobic, fermenting, eukaryotic parasite, we reasoned that *Giardia*'s unique anaerobic metabolism may disturb host gut ecology." (Eukaryotes are all organisms with cells that contain a nucleus, from microbes to plants to vertebrates. Prokaryotes, mostly bacteria, have much simpler cells, which lack a nucleus.)

Partly as a result of the unknown mechanism, there are few novel therapeutics, said Dawson. Moreover, even when treatment eradicates the parasite, GI symptoms may persist.

Provided by American Society for Microbiology

Citation: Giardiasis may be a disease of the ecology of the GI tract (2017, April 20) retrieved 10 April 2024 from

<https://medicalxpress.com/news/2017-04-giardiasis-disease-ecology-gi-tract.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.