

Negative sounds emphasized in burned-out brains

April 10 2017



Credit: George Hodan/public domain

Approximately every fourth working aged Finn experiences symptoms of burnout that include exhaustion, cynicism and reduced professional efficacy and often also difficulties in concentration and memory.

Licentiate in Psychology Laura Sokka from the University of Helsinki measured neural responses of 41 volunteers reporting a wide range of

[burnout](#) symptoms and their 26 controls during information processing and auditory tasks.

The results revealed that the participants experiencing burnout symptoms reacted faster to negative sounds of voice, and slower to positive utterances.

"It seems that those experiencing burnout symptoms are more sensitive to negative information," Sokka says.

The study is the first of its kind to study the EEG neural responses during demanding information processing tasks done by currently working people experiencing burnout symptoms.

In addition to listening tasks, the participants completed demanding attention and [memory tasks](#) simulating work life assignments. The tasks required quick decision-making in distractive surroundings and alternation between different [task](#) types.

Those experiencing mild burnout symptoms succeeded in the tasks well but their neural responses differed from their controls.

"Apparently, people with burnout symptoms struggle more in performing the tasks than the non-burnout controls. We observed decreased responses in the posterior scalp, and this decrement was compensated by increased responses in the frontal area," Sokka says.

In addition to neural changes, those experiencing severe burnout symptoms made more errors in the tasks.

"People with mild symptoms can cope with their workload quite long even though it's straining for the brain. When the symptoms worsen, they also start to make more errors," Sokka says.

The EEG measurements also revealed that the participants experiencing burnout symptoms did not react to sudden distractive noises as efficiently as their controls.

"There is a lot of discussion about how important it is to recover from stress and brain research supports this notion. We also know that long-lasting stress is a risk factor in many mental and physical illnesses, therefore, the current situation where every fourth has difficulties coping is not sustainable," Sokka says.

The tasks and measurements were repeated on the participants this spring and a follow-up study will later give evidence on how permanent the [neural response](#) changes are.

Provided by University of Helsinki

Citation: Negative sounds emphasized in burned-out brains (2017, April 10) retrieved 7 May 2024 from <https://medicalxpress.com/news/2017-04-negative-emphasized-burned-out-brains.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--