

# Man moves paralyzed legs using device that stimulates spinal cord

April 3 2017

---



Credit: Mayo Clinic

Mayo Clinic researchers used electrical stimulation on the spinal cord and intense physical therapy to help a man intentionally move his paralyzed legs, stand and make steplike motions for the first time in three years.

The case, the result of collaboration with UCLA researchers, appears

today in *Mayo Clinic Proceedings*. Researchers say these results offer further evidence that a combination of this technology and rehabilitation may help [patients](#) with [spinal cord injuries](#) regain control over previously paralyzed movements, such as steplike actions, balance control and standing.

"We're really excited, because our results went beyond our expectations," says neurosurgeon Kendall Lee, M.D., Ph.D., principal investigator and director of Mayo Clinic's Neural Engineering Laboratory. "These are initial findings, but the patient is continuing to make progress."

The 26-year-old patient injured his spinal cord at the sixth thoracic vertebrae in the middle of his back three years earlier. He was diagnosed with a motor complete spinal cord injury, meaning he could not move or feel anything below the middle of his torso.

The study started with the patient going through 22 weeks of physical therapy. He had three training sessions a week to prepare his muscles for attempting tasks during [spinal cord stimulation](#). He was tested for changes regularly. Some results led researchers to characterize his injury further as discomplete, suggesting dormant connections across his injury may remain.

Following physical therapy, he underwent surgery to implant an electrode in the epidural space near the spinal cord below the injured area. The electrode is connected to a computer-controlled device under the skin in the patient's abdomen. This device, for which Mayo Clinic received permission from the U.S. Food and Drug Administration for off-label use, sends electrical current to the spinal cord, enabling the patient to create movement.

After a three-week recovery period from surgery, the patient resumed

[physical therapy](#) with stimulation settings adjusted to enable movements. In the first two weeks, he intentionally was able to:

- Control his muscles while lying on his side, resulting in leg movements
  - Make steplike motions while lying on his side and standing with partial support
  - Stand independently using his arms on support bars for balance
- Intentional, or volitional, movement means the patient's brain is sending a signal to motor neurons in his spinal cord to move his legs purposefully.

"This has really set the tone for our post-surgical rehabilitation - trying to use that function the patient recovered to drive even more return of abilities," says Kristin Zhao, Ph.D., co-principal investigator and director of Mayo Clinic's Assistive and Restorative Technology Laboratory.

The Mayo researchers worked closely with the team of V. Reggie Edgerton, Ph.D., at UCLA on this study, which replicates earlier research done at the University of Louisville. The Mayo study marks the first time a patient intentionally controlled previously paralyzed functions within the first two weeks of stimulation.

The data suggest that people with discomplete [spinal cord](#) injuries may be candidates for epidural stimulation therapy. However, more research is needed into how a discomplete injury contributes to recovering function.

Teams from Mayo Clinic's departments of Neurosurgery and Physical Medicine and Rehabilitation, and the Division of Engineering collaborated on this project.

"While these are early results, it speaks to how Mayo Clinic researchers

relentlessly pursue discoveries and innovative solutions that address the unmet needs of patients," says Gregory Gores, M.D., executive dean of research, Mayo Clinic. "These teams highlight Mayo Clinic's unique culture of collaboration, which brings together scientists and physician experts who work side by side to accelerate scientific discoveries into critical advances for patient care."

**More information:** Peter J. Grahn et al. Enabling Task-Specific Volitional Motor Functions via Spinal Cord Neuromodulation in a Human With Paraplegia, *Mayo Clinic Proceedings* (2017). DOI: [10.1016/j.mayocp.2017.02.014](https://doi.org/10.1016/j.mayocp.2017.02.014)

Provided by Mayo Clinic

Citation: Man moves paralyzed legs using device that stimulates spinal cord (2017, April 3) retrieved 30 April 2024 from <https://medicalxpress.com/news/2017-04-paralyzed-legs-device-spinal-cord.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--