

# Peptide acts as mediator for learning

April 7 2017

---

The ability of the brain to respond and adapt to changes is scientifically called brain plasticity. This ability is the basis of all learning processes. New neurons, which can still be generated in the adult brain in specific areas, are instrumental in this process.

"But until now it has been unknown which molecular processes translate environmental changes into the production of new neurons," said Hannah Monyer, who leads the Cooperational Division of Clinical Neurobiology of the German Cancer Research Center (Deutsches Krebsforschungszentrum, DKFZ) and Heidelberg University Hospital. "In our current study, we have found a key mediator in this process for the first time."

Monyer and her team have revealed in their current study that a small peptide called DBI (short for: diazepam binding inhibitor) is the crucial mediator in this process. The peptide was first identified because it binds to the receptor for a chemical messenger in the [brain](#) called GABA, where it replaces a drug called Diazepam (Valium).

Recently, Monyer and colleagues already reported that DBI promotes the development of new neurons in an area of the brain called subventricular zone. This brain area is responsible for the supply of new nerve [cells](#) in the olfactory system, which is particularly sophisticated in rodents. In her present work, the Heidelberg neuroscientist shows that DBI has the same function in the hippocampus - the portion of the brain where memory formation and learning are located.

New neurons that form in the hippocampus improve orientation and learning capacity in the animals. Numerous research studies have already provided proof that in mice physical activity or variations in their environment stimulate neurogenesis in the hippocampus.

The researchers in Monyer's team used various genetic methods to turn off the DBI gene in this brain region in mice or, alternatively, to strongly boost it. When DBI was absent, the numbers of neural stem cells in the hippocampus declined. An oversupply of the peptide caused the opposite to happen, i.e., the investigators found more neural stem and progenitor cells.

Equipping the cages with toys is an established method to stimulate the generation of new neurons in the hippocampus in rodents. However, in mice whose DBI gene had been silenced using molecular-biological tricks, the stimulating environment failed to have an effect: The quantities of neural stem cells could not be increased.

DBI exerts its effect by binding to the receptor for the chemical messenger GABA in the [neural stem cells](#), thus acting as a molecular antagonist of this neurotransmitter. "GABA is responsible for keeping the stem cells dormant in their niche without dividing," explained Monyer. "When DBI enters the scene, they start proliferating, thus enlarging the stem cell pool that is available as a reservoir for young neurons. In DBI, we seem to have found the key mediator. The peptide suppresses the effect of GABA and thus links the environmental stimuli to the production of new [neurons](#) that are required for learning."

**More information:** Ionut Dumitru et al, Diazepam Binding Inhibitor Promotes Stem Cell Expansion Controlling Environment-Dependent Neurogenesis, *Neuron* (2017). [DOI: 10.1016/j.neuron.2017.03.003](https://doi.org/10.1016/j.neuron.2017.03.003)

Provided by German Cancer Research Center

Citation: Peptide acts as mediator for learning (2017, April 7) retrieved 27 April 2024 from <https://medicalxpress.com/news/2017-04-peptide.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.