

A quarter of nursing home residents are colonized with drug-resistant bacteria

April 27 2017

The significant presence of multidrug-resistant gram-negative bacteria (MDR-GNB), such as *E. coli*, among nursing home residents demonstrates the need for heightened infection control prevention and control measures in nursing homes, according to a meta-analysis published in the May issue of the *American Journal of Infection Control*, the official journal of the Association for Professionals in Infection Control and Epidemiology (APIC).

The systematic literature review and meta-analysis, conducted by Sainfer Aliyu, MPhil, MEd, MHPM, BSN, RN, and others at the Columbia University School of Nursing, found the prevalence of MDR-GNB colonization among sampled nursing home residents ranged from 11.2 percent to 59.1 percent, with a pooled average of 27 percent. Researchers also found that nine of the 12 studies involved identified specific factors that are associated with increased MDR-GNB colonization risk, including advanced age, gender, comorbid chronic diseases, history of recurrent hospitalization, increased interaction with healthcare workers, frequent antimicrobial exposure, delayed initiation of effective antibiotic therapy, presence of medical devices, decreased functional status, advanced dementia, nonambulatory status, fecal incontinence, severe sepsis present on admission, and residency in a long-term care facility.

The study is thought to be the first systematic review and meta-analysis of its kind, and should raise concerns among [infection control](#) professionals and nursing home facilities.

"This study underscores the importance of having strong infection prevention programs in all nursing homes and long-term care facilities," said 2017 APIC President Linda Greene, RN, MPS, CIC, FAPIC.

"Understanding the dynamics and cause of MDR-GNB transmission is crucial to identifying effective infection control strategies specific to these settings."

Nursing home residents are especially vulnerable to infections due to multiple comorbidities, advanced age, and immune dysfunction. In addition, living in a nursing [home](#) is itself considered a risk factor, as frequent transfers from nursing homes to acute care contribute to the influx of pathogens into hospital settings. Prevention and management of MDR-GNB in nursing homes are complicated and require extensive infection control resources due to challenges common to this setting such as understaffing, fewer resources, insufficient training, and inadequate surveillance.

"Identifying which patients are most prone to an increased risk of MDR-GNB will enable infection preventionists to tailor efforts and stem future contaminations," wrote Aliyu, et al. "The results of our study suggest that there is much more to be done with regard to [infection](#) prevention within [nursing homes](#), and that increased measures must be taken with elderly patients in regard to MDR-GNB colonization."

According to the Centers for Disease Control and Prevention, the percentage of gram-negative bacteria that are resistant to drugs is increasing. MDR-GNB cause serious infections in healthcare settings including pneumonia, bloodstream infections, wound or surgical site infections, and meningitis. They are particularly worrisome because they are becoming resistant to nearly all drugs that would be considered for treatment, with fewer novel antibacterial agents being developed.

More information: Sainfer Aliyu et al, Prevalence of multidrug-

resistant gram-negative bacteria among nursing home residents: A systematic review and meta-analysis, *American Journal of Infection Control* (2017). [DOI: 10.1016/j.ajic.2017.01.022](https://doi.org/10.1016/j.ajic.2017.01.022)

Provided by Elsevier

Citation: A quarter of nursing home residents are colonized with drug-resistant bacteria (2017, April 27) retrieved 5 May 2024 from <https://medicalxpress.com/news/2017-04-quarter-nursing-home-residents-colonized.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--