

Findings support role of vascular disease in development of Alzheimer's disease

April 11 2017



Diagram of the brain of a person with Alzheimer's Disease. Credit: Wikipedia/public domain.

Among adults who entered a study more than 25 years ago, an increasing number of midlife vascular risk factors, such as obesity, high blood pressure, diabetes, high cholesterol and smoking, were associated with elevated levels of brain amyloid (protein fragments linked to Alzheimer disease) later in life, according to a study published by *JAMA*.

Midlife vascular risk factors have been associated with late-life dementia. Whether these risk factors directly contribute to brain amyloid deposition is less well understood. Rebecca F. Gottesman, M.D., Ph.D., of the Johns Hopkins University School of Medicine, Baltimore, and

colleagues examined data from 346 participants without dementia at study entry who have been evaluated for vascular risk factors and markers since 1987-1989 and with PET scans in 2011-2013 as part of the Atherosclerosis Risk in Communities (ARIC)-PET Amyloid Imaging Study. Positron emission tomography image analysis was completed in 2015. Vascular risk factors at ARIC study entry (age 45-64 years; risk factors included [body mass index](#) 30 or greater, current smoking, hypertension, diabetes, and total cholesterol 200 mg/dL or greater) were evaluated in models that included age, sex, race, APOE genotype, and educational level.

The availability of imaging biomarkers for brain amyloid allows the study of individuals before the development of dementia and thereby allows consideration of the relative contributions of vascular disease and amyloid to cognition, as well as the contribution of vascular disease to amyloid deposition.

The researchers found that a cumulative number of midlife vascular risk factors were associated with elevated brain amyloid. Relationships between vascular risk factors and brain amyloid did not differ by race. The results were not supportive of a significant difference in association among people who were or were not carriers of an APOE ϵ 4 allele (a variant of a gene associated with increased risk for Alzheimer disease). Late-life vascular risk factors were not associated with late-life brain amyloid deposition.

"These data support the concept that midlife, but not late-life, exposure to these [vascular risk factors](#) is important for [amyloid deposition](#)," the authors write. "These findings are consistent with a role of vascular disease in the development of AD."

More information: *JAMA* (2017). [DOI: 10.1001/jama.2017.3090](https://doi.org/10.1001/jama.2017.3090)

Provided by The JAMA Network Journals

Citation: Findings support role of vascular disease in development of Alzheimer's disease (2017, April 11) retrieved 27 April 2024 from <https://medicalxpress.com/news/2017-04-role-vascular-disease-alzheimer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.