


Specific factors influence first-line biologic Rx for psoriasis

April 13 2017



(HealthDay)—The presence of psoriatic arthritis, patient weight,

registration country, employment status, and disease severity are the main factors influencing first-line biologic treatment selection for patients with psoriasis, according to a study published online April 7 in the *British Journal of Dermatology*.

Niall J. Davison, from the University of Manchester in the United Kingdom, and colleagues used data from the British Association of Dermatologists Biologic Interventions Register (January 2012 to December 2015) to determine the factors which influenced the probability of first-line biologic treatment selection for 3,040 individuals with psoriasis.

The researchers found presence of [psoriatic arthritis](#), patient weight, [employment status](#), country of registration, and baseline [disease severity](#) were factors affecting first-line biologic selection. Over time, prescribing behavior changed, possibly due to experiences and emerging evidence on treatment effectiveness and safety.

"[This study] provides baseline data to inform the evaluation of future potential changes that may impact prescribing behavior such as stratified medicine," the authors write.

Several authors disclosed financial ties to the pharmaceutical industry.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Specific factors influence first-line biologic Rx for psoriasis (2017, April 13) retrieved 27 April 2024 from <https://medicalxpress.com/news/2017-04-specific-factors-first-line-biologic-rx.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.