

Disfiguring eye symptoms diminish in Graves' eye disease drug trial

May 3 2017



Research led Terry J. Smith, M.D., and Raymond Douglas, M.D., Ph.D., while at the University of Michigan Kellogg Eye Center, shows success of teprotumumab in safely reducing disfiguring eye symptoms of Graves' eye disease. Credit: University of Michigan Kellogg Eye Center

The disease is characterized by inflammation and a buildup of tissue



around the eyes that cause them to bulge painfully from their sockets.

There has never been a safe and effective treatment for Graves' <u>eye</u> <u>disease</u>, also known as thyroid eye disease, for the one million Americans with the condition. Patients see inconsistent benefit from antiinflammatory medicines and complex orbital surgery helps give some a more natural appearance once the disease has run its course.

The results of a clinical trial led by the University of Michigan Kellogg Eye Center offers hope for those with moderate-to-severe active TED. Patients had a significant reduction in eye symptoms after treatment with teprotumumab, a study drug the U.S. Food and Administration called a "breakthrough therapy."

Most study <u>patients</u> in the trial who got the active drug instead of a placebo had reduced eye bulging, eye pain and swelling and improved quality of life, the study's primary and secondary end points.

For some, the therapeutic benefit was rapid, occurring within six weeks of intravenous infusions of the drug that blocks what researchers say is an <u>autoimmune attack</u> on muscles around the eyes.

Published in the *New England Journal of Medicine*, the clinical trial data represents 25 years of research led by the lead study author the Kellogg Eye Center's Terry J. Smith, M.D., an endocrinologist and Frederick G.L. Huetwell Professor of Ophthalmology and Visual Sciences.

TED is a rare condition associated with Graves' disease, an autoimmune disorder that causes an overproduction of thyroid hormone, or hyperthyroidism. Symptoms include swelling of the eyelids, constant stare, bulging eyes, eyelid retraction, optic neuropathy, double vision and other symptoms.



Approximately 25 to 50 percent of those with Graves' disease will develop TED, most of them women.

Previous findings in Smith's laboratory suggested that an insulin-like growth factor 1 receptor (IGF-1R) plays an important role in Graves' disease and other autoimmune diseases.

"Specifically, we have found that there is an unusually high presence of IGF-1 receptors on cells surrounding the eye in TED," says Smith, who is widely recognized for his groundbreaking research on treatments for TED and related autoimmune disorders.

Building on this discovery, an outgrowth of basic science research conducted at the University of California, Los Angeles, and then at the Kellogg Eye Center, Smith and study author Raymond Douglas, M.D., Ph.D., a Graves' eye disease specialist and renowned oculoplastics surgeon, initiated the study of thyroid-associated ophthalmopathy.

"The eyes are particularly susceptible to Graves' disease because the autoimmune attack often targets the eye muscles and connective tissue surrounding the eye," Douglas says. "This occurs as a consequence of these tissues containing proteins that are shared with the thyroid gland."

The study, lasting 24 weeks and involving 88 patients, was the largest multi-center Graves' eye disease trial. Patients were recruited between July 2, 2013 and Sept. 23, 2015, from 15 sites across the United States as well as Italy and Germany and the United Kingdom.

Study investigators examined the benefit of teprotumumab, a monoclonal antibody shown to inhibit IGF-1 function and was originally developed as a cancer drug.

Although teprotumumab failed to exhibit efficacy in patients with



cancer, the team believed it might be useful in interrupting the TED disease process.

At the end of the trial, 69 percent of study patients receiving infusions of teprotumumab once every three weeks had reduced eye bulging (proptosis) and increased quality of life compared to 20 percent in the placebo group.

Patients were randomly assigned to receive either a placebo or an active drug. Neither participants nor researchers knew which medicine was being used so they could describe what happened in the trial without bias.

Some participants had results after two injections at week six when, according to the study, a reported 43 percent of patients in the treatment group compared to 4 percent in the placebo group had a response.

Patients who responded to the treatment reached the study's primary end point of a reduction by 2 points or more in the severity of the disease, which was measured on a 7-point Clinical Activity Score, and reduction by 2 mm or more in the degree of proptosis. Secondary end points included better quality of life.

Those in the treatment group had a baseline Clinical Activity Score of 5.1. The marked reduction in proptosis is similar to the kind of results surgical patients would expect, authors note.

"The findings have implications for other autoimmune diseases in which IGF-1 reception has already been implicated such as rheumatoid arthritis," Smith says.

The most common side effect of treatment was nausea. Muscle spasms and diarrhea were also reported, particularly among patients with



gastrointestinal <u>disease</u>. The only drug-related adverse event was hyperglycemia in patients with diabetes; these events were controlled by adjusting medication for diabetes.

The results provide inroads for a new therapy to safely minimize the severity of the eye condition.

"The study is one of two pivotal <u>clinical trials</u> needed to achieve registration of teprotumumab as the first approved therapy to prevent TED symptoms," says David Madden, president and chief executive officer of River Vision Corporation, which has entered an agreement to manufacture the drug with CMC Biologics in Copenhagen, Denmark.

The drug cannot be prescribed currently for TED patients. A follow-up clinical trial is underway, but not recruiting participants, and is expected to be complete in 2018.

Provided by University of Michigan

Citation: Disfiguring eye symptoms diminish in Graves' eye disease drug trial (2017, May 3) retrieved 8 May 2024 from <u>https://medicalxpress.com/news/2017-05-disfiguring-eye-symptoms-diminish-graves.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.