

Early epigenetic switches associated with childhood bone health

May 10 2017

The health of children's bones could be determined before they are born, a new University of Southampton study has shown.

Researchers from the MRC Lifecourse Epidemiology Unit and the Institute of Developmental Sciences at the University of Southampton, as part of the Epigen Global Consortium, looked at whether [bone health](#) might be influenced by [epigenetic modifications](#) of DNA early in life.

The results, published in the *Journal of Bone and Mineral Research*, provide an insight into the early determinants of skeletal growth, and improve the understanding of how osteoporosis could be prevented in future generations.

There is growing evidence that whether genes are expressed or not (switched on or off) in particular human cells can change throughout life and can be affected by a range of environmental factors even before birth, such as their parents' health, diet and lifestyle before and during pregnancy. This switching on or off of genes is known as "epigenetic modification" and an important epigenetic mechanism is DNA methylation.

The Southampton researchers analysed the levels of DNA methylation in umbilical cord tissue of 669 babies born in the Southampton Women's Survey.

They compared the DNA methylation levels in the CDKN2A gene to the

[bone mass](#) of the child at four and six years of age, measured using DXA [bone](#) densitometry. They found that higher DNA methylation in particular parts of the CDKN2A gene, which is known to play a role in development and ageing, was associated with lower bone mass at four and six years.

Analysis showed that a 10 percent increase in methylation was associated with a decrease in total bone mass of around 4-9g at age four years. Further laboratory analysis showed that methylation of the CDKN2A region is important for the function and survival of [bone cells](#).

Nicholas Harvey, Professor of Rheumatology and Clinical Epidemiology at the University of Southampton, led the study with Dr Elizabeth Curtis, Wellcome Trust Clinical Research Fellow, and Dr Robert Murray, Postdoctoral Research Fellow, both also from the University.

He said: "The health of a child's bone when they are young can influence the risk of osteoporosis in older age. This study provides exciting insights into the role of epigenetics in bone health, and might allow us to more accurately predict an individual's future risk of osteoporosis. Our ongoing studies should enable us to work out whether interventions during pregnancy, for example vitamin D supplementation, will actually alter the epigenetic marks, and lead to improved bone [health](#) in the offspring."

Professor Cyrus Cooper, Director of the MRC Lifecourse Epidemiology Unit, said: "This major finding links our previous observations on maternal nutrition and lifestyle during pregnancy, with the later risk of musculoskeletal ageing in the offspring. It bears testimony to the value of large, well-maintained population cohorts, participants among whom are followed up for many years."

More information: Elizabeth M Curtis et al. Perinatal DNA

methylation atis associated with offspring bone mass: Findings from the Southampton Women's Survey, *Journal of Bone and Mineral Research* (2017). [DOI: 10.1002/jbmr.3153](https://doi.org/10.1002/jbmr.3153)

Provided by University of Southampton

Citation: Early epigenetic switches associated with childhood bone health (2017, May 10)
retrieved 19 April 2024 from
<https://medicalxpress.com/news/2017-05-early-epigenetic-childhood-bone-health.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.