

Early treatment for NSTEMI patients shows greater rate of survival

May 12 2017

An analysis of NSTEMI patients who undergo coronary revascularization within 24 hours of hospitalization showed an increased reduction in mortality, marking the first time this difference has been demonstrated. Results from "Outcomes of Early vs. Late Revascularization in Low and High-Risk Patients Hospitalized with Non-ST-Elevation Myocardial Infarction: The Atherosclerosis Risk in Communities (ARIC) Surveillance Study" were presented today as a late-breaking clinical trial at the Society for Cardiovascular Angiography and Interventions(SCAI) 2017 Scientific Sessions in New Orleans.

NSTEMI, or Non-ST-Elevation Myocardial Infarction, is a type of heart attack that occurs when an artery is only partially blocked as opposed to a STEMI, or ST-Elevation Myocardial Infarction, where there is a complete blockage of the artery, making it the more severe of the two types.

While current guidelines recommend early intervention—defined as less than 24 hours—for STEMI [patients](#), delayed or late strategy—defined as between 24-72 hours—for NSTEMI is considered reasonable, unless there are extenuating circumstances, such as refractory angina or other conditions that put the patient in a higher risk category.

However, the optimum time for coronary revascularization of NSTEMI patients is under debate, according to Sameer Arora, MD, of the University of North Carolina at Chapel Hill, Division of Cardiology and the study's lead investigator. Evidence for current recommendations is

based on clinical trials in controlled settings and selected patients.

Arora and his collaborators looked at data from the ARIC Community Surveillance Study, a large, ongoing investigation that began in 1987 involving 21 hospitals in four states: Maryland, Minnesota, Mississippi, and North Carolina. Arora analyzed data from hospitalized NSTEMI patients undergoing [coronary revascularization](#), and classified them as low or high risk, based on accepted risk scores. The survival benefit of an early revascularization(

Citation: Early treatment for NSTEMI patients shows greater rate of survival (2017, May 12) retrieved 4 May 2024 from

<https://medicalxpress.com/news/2017-05-early-treatment-nstemi-patients-greater.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.