

Infertility in men could point to more serious health problems later in life

May 16 2017, by Aleksander Giwercman And Yvonne Lundberg
Giwercman



Credit: AI-generated image ([disclaimer](#))

Poor sperm quality affects about [one in ten men](#) and may lead to fertility problems. These men also have an increased risk of [developing testicular cancer](#), which is the most common malignant disease of young males. And, even if they don't develop testicular cancer, men with poor sperm

quality [tend to die younger](#) than men who don't have fertility problems.

Couples who can't achieve pregnancy usually go to fertility clinics for treatment. At these clinics, emphasis is put on deciding whether the couple needs assisted reproduction or not, and, if so, to choose between different methods (such as [IVF](#), [IUI](#), or [ICSI](#)) for doing this. In most cases, these treatments lead to pregnancy and a live birth. So the problem seems to be solved. But if infertility is an early symptom of an underlying disease in the man, fertility clinics won't pick it up.

Missed opportunity

Testicular cancer is easy to detect. In men seeking treatment for [fertility problems](#), a simple ultrasound scan of the testes can reveal early cancer, so a life-threatening tumour can be prevented. If detected, [95% of all cases can be cured](#). But, unfortunately, testicular ultrasound scans are rarely performed at fertility clinics as the focus tends to be on sperm numbers and which method of assisted reproduction to use.

And [testicular cancer](#) is not the only threat to young infertile men's health. [Serious health problems](#), such as metabolic syndrome ([high blood pressure](#), [high blood sugar](#) and obesity), type 2 diabetes and loss of bone mass are also much more common conditions among infertile men. These disorders are possible to prevent, but if left untreated often lead to premature death.

A possible culprit

At Lund University in Malmö, Sweden, we have – together with other research groups – made a number of studies focusing on the link between male fertility problems and subsequent risk of serious diseases. We cannot yet explain the causes, but testosterone deficiency is a strong

candidate. My research team found that [30% of all men with impaired semen quality](#) have [low testosterone levels](#). And men totally lacking the hormone have early signs of diabetes and bone loss.

We recently conducted [a study](#) in which we investigated almost 4,000 men below the age of 50 and who had had their testosterone measured 25 years ago. We found that the risk of dying at a young age was doubled among those with low testosterone levels compared with men with normal levels of this hormone.

Although testosterone treatment may not necessarily be the best preventive measure, these findings makes it possible to identify men at high risk so that they can be advised about lifestyle changes, such as losing weight or quitting smoking – lifestyle changes that will help reduce the risk of developing type 2 diabetes, cardiovascular disease and osteoporosis.

A relatively high proportion of men get in touch with their doctor about infertility [problems](#) and, as they represent a high-risk group for some of the most common diseases occurring later in life, perhaps it is time to change the routines for managing them. With the knowledge we now have regarding these men's health, the least we can demand from doctors is to identify those who are at risk of serious diseases after they have become fathers. This is cheap and only requires simple tests. It is no longer enough to just evaluate the number of sperm.

This article was originally published on [The Conversation](#). Read the [original article](#).

Provided by The Conversation

Citation: Infertility in men could point to more serious health problems later in life (2017, May

16) retrieved 2 May 2024 from <https://medicalxpress.com/news/2017-05-infertility-men-health-problems-life.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.