

Finding traces of memory processing during sleep

May 17 2017



Credit: Vera Kratochvil/public domain

Sleep helps us to retain the information that we have learned during the

day. We know from animal experiments that new memories are reactivated during sleep. The brain replays previous experience while we sleep – and this replay strengthens memories overnight. Up to now, it was hard to demonstrate such a reactivation in humans, because the activity of individual neurons cannot be observed and most memories will activate entire networks of brain regions. Scientists from Tübingen have now applied new statistical pattern detection methods from the field of machine learning to get around these problems.

The researchers recorded [brain](#) activity during sleep with electroencephalography (EEG). They measured electrical brain activity in participants who had previously studied different kinds of pictures under varied learning conditions. They then trained a support vector machine – a computer-ized algorithm for pattern recognition – to distinguish these different learning conditions. The algorithm was then applied to new test subjects, and was able to predict which kinds of pictures they had studied before sleep – based only on their electrical brain activity during sleep. Learning those images must thus have impacted the sleeping brain's activity.

In their study, which has been published in *Nature Communications*, the neuroscientists show that humans also reprocess recent memories in sleep. They found signs of such a reprocessing in deep [slow-wave sleep](#) as well as in REM sleep. "The stronger the signs of reprocessing in sleep, the better our subjects remembered the material in the morning. Thus, our study provides initial evidence that sleep-dependent processing of memories actually leads to their stabilization", explains Dr. Monika Schönauer, who has implemented the study and analyzed the data together with her colleague Sarah Alizadeh. Most interestingly, the influence of reprocessing on memory retention was found in slow-wave sleep, but not in REM sleep. "Slow-wave sleep and REM sleep both seem to participate in [memory](#) processing. However, they must have different functions", concludes Schönauer from these findings. Now it is

particularly important to gain an understanding of the function of REM sleep, she says.

A major part of the research project was finding a method that could identify precisely those fluctuations of brain electrical activity in sleep which were caused by previous learning. "We had to find a way to detect tiny, learning-related patterns in the vast expanse of brain [activity](#) during the night", says Dr. Steffen Gais, who was the leading scientist. With the help of multivariate [pattern recognition](#) algorithms, the researchers were able to extract specific information about previously learned material from human [electrical brain activity](#). Gais is confident with regard to future studies: "Pattern recognition is a highly sensitive method, which is becoming more and more widely adopted in the life sciences. I am certain that it will enable breakthroughs especially in cognitive neurosciences, where it will allow us to investigate previously hidden processes like dreams and spontaneous thought processes."

More information: M. Schönauer et al. Decoding material-specific memory reprocessing during sleep in humans, *Nature Communications* (2017). [DOI: 10.1038/NCOMMS15404](https://doi.org/10.1038/NCOMMS15404)

Provided by Universitaet Tübingen

Citation: Finding traces of memory processing during sleep (2017, May 17) retrieved 25 April 2024 from <https://medicalxpress.com/news/2017-05-memory.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--