

Hand osteoarthritis is a common condition

May 4 2017

A new study estimates that the lifetime risk of symptomatic hand osteoarthritis is 40%, and nearly one in two women and one in four men will develop the condition, which affects hand strength and function and causes disability in activities of daily living.

Race-specific estimates are 41% among whites and 29% among blacks. Also, the [lifetime risk](#) among individuals with obesity is 47%, which is 11 percentage points higher than those without obesity.

These findings indicate that hand osteoarthritis is very common. In comparison, prior studies have estimated lifetime risks for symptomatic knee and [hip osteoarthritis](#) to be 45% and 25%, respectively.

"These findings demonstrate the substantial burden of symptomatic hand osteoarthritis overall and in subgroups. Increased use of public health and clinical interventions is needed to address its impact," wrote the authors of the *Arthritis & Rheumatology* study.

More information: Jin Qin et al, Lifetime Risk of Symptomatic Hand Osteoarthritis: The Johnston County Osteoarthritis Project, *Arthritis & Rheumatology* (2017). [DOI: 10.1002/art.40097](https://doi.org/10.1002/art.40097)

Provided by Wiley

Citation: Hand osteoarthritis is a common condition (2017, May 4) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2017-05-osteoarthritis-common-condition.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.