

Physical activity rather than sedentary behaviour affects the risk of mortality

May 4 2017

It is not the amount of time spent sitting still that matters. Instead, the extent of physical activity is essential in reducing the risk of elderly women developing cardiovascular disease, as shown in a new Örebro study published in *PLOS ONE*.

"We have studied women over 65, as they are among the least active groups of the population, and run a high risk of developing [cardiovascular disease](#)," says Fawzi Kadi, Professor at Örebro University.

A [sedentary lifestyle](#) is associated with health risks such as diabetes and cardiovascular disease. But the Örebro study shows that it is the daily amount of [physical activity](#) that is crucial to a person's health.

"The study shows how important it is to encourage more physical activity. We are not talking slow everyday pace, but at least one brisk walk or other physical activity requiring some exertion," says Andreas Nilsson, researcher at Örebro University.

120 women took part in the Örebro study. After a medical examination, their physical activity was measured using an accelerometer over the course of one week.

The results may well apply to other groups, since they are in line with a meta-analysis of previous research based on a million adult men and women, which indicated that physical activity rather than sedentary

behaviour affects the risk of mortality.

"Our study points in the same direction – that the [negative health effects](#) of a sedentary lifestyle decrease with the extent of physical activity," says Fawzi Kadi.

This means that if one person is jogging while another is only doing less strenuous activities, the first person runs a lower risk of developing cardiovascular disease than the second – regardless of the extent of their [sedentary activities](#).

"Getting up once in awhile is naturally a good thing, but doing more exercise is better for our health," concludes Andreas Nilsson.

More information: Andreas Nilsson et al. Physical activity and not sedentary time per se influences on clustered metabolic risk in elderly community-dwelling women, *PLOS ONE* (2017). [DOI: 10.1371/journal.pone.0175496](#)

Provided by Örebro Universitet

Citation: Physical activity rather than sedentary behaviour affects the risk of mortality (2017, May 4) retrieved 19 April 2024 from <https://medicalxpress.com/news/2017-05-physical-sedentary-behaviour-affects-mortality.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|