

Study spotlights Indigenous children and serious burns injuries

May 15 2017, by Gabrielle Dunlevy



Credit: AI-generated image (disclaimer)

More Indigenous children are going to hospital in NSW with serious burns than non-Indigenous children and they are less likely to be treated in a hospital in a paediatric burns unit, despite needing more intensive treatment and a longer stay.



A study led by the UNSW Centre for Big Data Research in Health and supported by researchers at The George Institute for Global Health investigated the differences in <u>burn</u> injuries in children, examining the cause of the injury, its location on the body, the total body surface affected (% TBSA), burn depth and length of <u>hospital stay</u>.

The proportion of Indigenous children with burns who presented with injuries affecting more than 10% TBSA was greater and the hospital stay was usually almost three days longer than non-Indigenous children.

A smaller proportion of Indigenous children with burns were treated in a hospital with a paediatric tertiary burns unit, fitting with previous studies that have shown Indigenous Australians experience inequities in access to medical services.

The first author on the paper published today in the *Medical Journal of Australia*, Holger Möller, says the higher proportion of Indigenous children presenting with burns affecting more than 10% TBSA is of particular concern.

"Burns can be among the most devastating of child injuries and can result in long-term physical and psychological impairment thus affecting the child's development and future life," he said.

"We could not assess the longer term outcomes of burn injury in this study and to date little is known about the long term outcomes, the post-discharge care, and the impact of care on functional outcomes in Aboriginal children."

Scalds were the leading cause of burn <u>injury</u> to both Indigenous (47%) and non-Indigenous children (62%). There was a higher proportion of flame burns in Indigenous children, which may be partially explained by the higher number of Indigenous children living in rural and remote



areas where there are more outdoor fires.

The study involved population-based cohort analysis of linked hospital and mortality data for 2000–2014, with 35,749 Indigenous and 1,088,938 non-Indigenous children aged under 13 years as participants.

The study was done with researchers from Neuroscience Research Australia, Australian National University, The George Institute for Global Health, Flinders University and the University of Wollongong.

Study author, Professor Rebecca Ivers, director of the Injury Division at The George Institute, is currently exploring the care of Aboriginal children with burns through a cohort study in Queensland, NSW, South Australia and the Northern Territory.

The study is following children for at least two years post-burn in order to understand the impact and cost of burns. A roundtable is being planned for 2018 to develop a new model of care. The study utilises Indigenous research methodologies and three Aboriginal PhD students are working on the study.

"The study builds on previous understanding about burns in Aboriginal and Torres Strait Islander children but rather than just measuring inequality, will result in a new transformative model of care that meets the needs of patients and caregivers," Professor Ivers said.

More information: Holger Möller et al. Indigenous and non-Indigenous Australian children hospitalised for burn injuries: a population data linkage study, *The Medical Journal of Australia* (2017). DOI: 10.5694/mja16.00213



Provided by University of New South Wales

Citation: Study spotlights Indigenous children and serious burns injuries (2017, May 15) retrieved 19 April 2024 from

https://medicalxpress.com/news/2017-05-spotlights-indigenous-children-injuries.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.