


Stem-cell transplants show limited benefit for double-hit lymphoma patients in remission

May 15 2017



Hodgkin lymphoma, nodular lymphocyte predominant (high-power view) Credit: Gabriel Caponetti, MD./Wikipedia/CC BY-SA 3.0

Patients with double hit lymphoma (DHL) who undergo autologous stem-cell transplantation (autoSCT) after achieving remission are not more

likely to remain in remission or live longer than patients who do not undergo autoSCT, according to a new analysis from the Perelman School of Medicine and the Abramson Cancer Center of the University of Pennsylvania. The study looked at long term outcomes for patients who achieved remission and, in most cases, found no clear benefit to the transplant, except potentially in patients who received standard front-line chemotherapy, who were less likely to remain in remission than those patients receiving intensive front-line chemotherapy. The findings are published this month in the *Journal of Clinical Oncology*.

DHL is a form of aggressive B cell non-Hodgkin lymphoma characterized by genetic alterations that drive the lymphoma's growth. This variant is associated with a poor prognosis as compared to other forms of aggressive B cell lymphomas, as patients with this disease survive only an average of two years after diagnosis. Relapses of this disease are almost always fatal, meaning that keeping patients in remission is crucial.

"A major dilemma for oncologists who treat this disease was whether or not to recommend the potentially harmful therapy of autoSCT to patients with this disease a strategy to help keep them in remission," said Daniel J. Landsburg, MD, an assistant professor of Hematology Oncology at Penn and the study's lead author.

Landsburg and his team looked at data on 159 patients from 19 different academic medical centers across the United States. Patients were diagnosed between 2006 and 2015, and all achieved remission following intensive front-line chemotherapy or the standard chemotherapy regimen containing rituximab plus cyclophosphamide, doxorubicin, vincristine, and prednisone (R-CHOP). Of the total patients, 62 underwent an autoSCT, while 97 did not. Landsburg noted that there were no significant differences between the patient groups at baseline.

"Our result is not explained by differences in patients' overall health or disease features," Landsburg said. "The transplant and non-transplant arms of this study were very well-matched."

Overall, 80 percent of the patients were still in remission three years later, and 87 percent were still alive. When researchers broke the patients into two groups, autoSCT and no autoSCT, they found 89 percent of autoSCT patients were still in remission at three years, as were 75 percent of patients who did not receive an autoSCT. Also at three years, 91 percent of autoSCT patients were still alive, compared to 85 percent of non-autoSCT patients. None of these differences were found to be statistically significant.

"Once these patients achieve remission, the data show they are likely to stay in remission," Landsburg said. "In the absence of a large randomized controlled trial, which would be very challenging to carry out in this case, this is the best evidence we have, and it shows there's no clear benefit to these patients undergoing autoSCT."

Landsburg did point to one exception in the data, and that was in patients who underwent RCHOP, the standard front-line chemotherapy regimen. Just 56 percent of them were still in remission at three years, far lower than patients who received the more intensive front-line therapies.

"Even if patients do go into remission with RCHOP, it appears to be less durable, so in these cases, going forward with autoSCT may still make sense," Landsburg said.

Landsburg says the next step will be to study features of patients who don't go into [remission](#) in order to understand why their disease is resistant to therapy and if that can be overcome with different treatment strategies. He says it's also important to try to find more effective therapies for DHL patients who relapse.

More information: Daniel J. Landsburg et al, Outcomes of Patients With Double-Hit Lymphoma Who Achieve First Complete Remission, *Journal of Clinical Oncology* (2017). [DOI: 10.1200/JCO.2017.72.2157](https://doi.org/10.1200/JCO.2017.72.2157)

Provided by Perelman School of Medicine at the University of Pennsylvania

Citation: Stem-cell transplants show limited benefit for double-hit lymphoma patients in remission (2017, May 15) retrieved 11 May 2024 from <https://medicalxpress.com/news/2017-05-stem-cell-transplants-limited-benefit-double-hit.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.