

May 23 2017

Strategic brain training positively affects neural connectivity for individuals with TBI



A thresholded Z-statistical maps and anatomical views for between-group differences in resting-state functional connectivity changes over time and at TP3 (|Z|>1.96, pNBS

A recent study from the Center for BrainHealth at The University of Texas at Dallas shows that a certain type of instructor-led brain training protocol can stimulate structural changes in the brain and neural connections even years after a traumatic brain injury (TBI).

The findings, published in *Brain and Behavior*, further suggest that changes in <u>cortical thickness</u> and neural network connectivity may prove an effective way to quantitatively measure treatment efficacy, an ability that has not existed until now. Building upon previous research, the study challenges the widely held belief that recovery from a TBI is limited to two years after an injury.



"A TBI disrupts brain structure. These brain changes can interfere with brain network communication and the cognitive functions those networks support," said Dr. Kihwan Han, research scientist at the Center for BrainHealth and lead author of the study.

"For people with chronic TBI, they may have trouble with daily tasks such as creating shopping lists and resolving conflicts with others for many years after the injury. These findings provide hope for people who thought, 'This is as good as my recovery is going to get' and for the medical community who have yet to find a way to objectively measure a patient's recovery," he said.

The study included 60 adults with TBI symptoms lasting an average of eight years. Participants were randomly placed into one of two cognitive <u>training</u> groups: strategy-based training or knowledge-based training. Over an eight-week period, the strategy-based training group learned strategies to improve attention and reasoning. The knowledge-based training group learned information about the structure and function of the brain as well as the effects of sleep and exercise on brain performance.

Magnetic resonance imaging measured cortical thickness and resting-state functional connectivity (rsFC) before training, after training and three months post-training. Previous studies have shown that cortical thickness and rsFC can be potential markers for training-induced brain changes.

Individuals in the strategy-based reasoning training showed a greater change in cortical thickness and connectivity compared to individuals who received the knowledge-based training. Changes in cortical thickness and functional connectivity also correlated to an individual's ability to switch between tasks quickly and consistently to achieve a specific goal.

"People who showed the greatest change in cortical thickness and connectivity, showed the greatest performance increases in our cognitive tasks. Perhaps future studies could investigate the added benefit of <u>brain</u> stimulation treatments in combination with cognitive training for individuals with chronic TBI who experience problems with attention, memory or executive functions," Han said.

More information: Kihwan Han et al, Strategy-based reasoning training



modulates cortical thickness and resting-state functional connectivity in adults with chronic traumatic brain injury, *Brain and Behavior* (2017). <u>DOI:</u> <u>10.1002/brb3.687</u>

Provided by Center for BrainHealth

Citation: Strategic brain training positively affects neural connectivity for individuals with TBI (2017, May 23) retrieved 1 May 2024 from <u>https://medicalxpress.com/news/2017-05-strategic-brain-positively-affects-neural.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.