

A new test to detect anti-phospholipid antibody syndrome

June 14 2017

In autoimmune diseases, the immune system wrongly produces antibodies that attack the patient's own cells. One of these diseases, anti-phospholipid antibody syndrome (APS), is still poorly understood, even though it can have serious consequences. APS is caused by antibodies circulating in the blood plasma that are directed against a protein, which increases the blood's tendency to form clots. This can lead to a range of vascular problems such as venous thromboses, strokes or repeated miscarriages.

Although the prevalence of APS is very difficult to assess, it is likely to affect around 0.5% of the general population. Diagnosing the disease is a complicated affair. The most commonly used test has a number of problems in terms of variability, specificity and sensitivity. However, researchers at the University of Geneva (UNIGE), Switzerland, and the Geneva University Hospitals (HUG) have succeeded in identifying the exact site where the anti-phospholipid [antibodies](#) attach themselves. This means a more accurate and standardized diagnostic test can now be devised—an undeniable improvement for patients. These results are published in the journal *Haematologica*.

In people suffering from APS, antibodies called anti-B2GP1 attach themselves to elements found on the surfaces of certain cells, particularly those of the blood vessels and placenta. They bind themselves to receptors located on the cell membrane, generating a signal that produces the pro-inflammatory and pro-thrombotic factors that cause vascular incidents. By identifying the exact location where

these antibodies bind, the research team at UNIGE and HUG have clarified how they function. Karim Brandt, a researcher at the UNIGE Faculty of medicine, explains the importance of this discovery: "The current diagnostic tests use the entire protein, which reduces its specificity and leads to standardization issues. Consequently, two tests are required at an interval of 12 weeks after a thrombotic episode or following one or more miscarriages. Our new test specifically targets this pathogenic antibody, with rapid and more accurate results."

The researchers managed to isolate a small part of the membrane protein referred to as a motif. Motifs are recognized by the antibody, which then binds to it, like a key in a lock. In this instance, the key can open several locks that correspond to the proteins found on the surfaces of the cells and induce the [pathogenic effects](#). It is the only protein in all the human proteome to have five of these motifs; it therefore has as many potential binding points for the pathogenic antibody.

Better diagnosis and better treatment

APS is usually treated with oral anticoagulants such as low-molecular-weight heparin and aspirin, long-term treatments that are not without side effects, and that must be used with caution by pregnant women. Moreover, treatment becomes burdensome in patients suffering from the most severe form of the disease, called catastrophic APS. Karim Brandt stresses that the researchers are also focusing their studies in this direction: "Our breakthrough could also give rise to a targeted treatment that would neutralize specific pathogenic antibodies, reducing not just their actions but also the side effects associated with the current treatment. It would involve injecting the protein motif we have identified into a patient's circulatory system so that it explicitly binds itself to the pathogenic antibody and prevents it from causing harm."

For the time being, the diagnostic test needs to be optimized for

prototypes to be developed. To ensure its validity, the researchers will reanalyze hundreds of samples already tested with the old method and compare results.

More information: Philippe de Moerloose et al. Patient-derived anti- β 2GP1 antibodies recognize a peptide motif pattern and not a specific sequence of residues, *Haematologica* (2017). DOI: [10.3324/haematol.2017.170381](https://doi.org/10.3324/haematol.2017.170381)

Provided by University of Geneva

Citation: A new test to detect anti-phospholipid antibody syndrome (2017, June 14) retrieved 2 May 2024 from <https://medicalxpress.com/news/2017-06-anti-phospholipid-antibody-syndrome.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--