

## **Cardiac arrest on plane journeys: New guidelines proposed**

June 5 2017



Credit: CC0 Public Domain

New guidelines to deal with the in-flight emergency of cardiac arrest in a passenger or crew member are being proposed at this year's Euroanaesthesia meeting in Geneva (3-5 June). The task force that has



created the guidelines is led by Professor Jochen Hinkelbein, University of Cologne, Germany and President of the German Society for Aerospace Medicine (DGLRM).

Approximately 3 billion people worldwide travelled by commercial air transport in 2016. A calculation based on the number of passengers transported shows that between 1 out of 14,000 to 1 out of 50,000 passengers will experience acute medical problems during a flight. While cardiac arrest accounts for just 0.3% of all in-flight medical emergencies, it is responsible for 86% of in-flight events resulting in death. Recently, the Star Wars actress Carrie Fisher died following a cardiac arrest she suffered on a plane journey to Los Angeles.

Despite this high fatality rate of in-flight cardiac arrest (IFCA), so far, no guideline for IFCA exists that provides specific treatment recommendations. Thus, a DGLRM <u>task force</u> was created to develop a guideline for the treatment of in-flight cardiac arrest based on clinical and investigational expertise in this area.

Several main recommendations have been developed, which include:

- Emergency equipment provided and its location should be mentioned in the pre-flight safety announcement
- An electrocardiogram (ECG) should be available for patients with cardiac arrest; automated external defibrillators (AEDs) have this function available and many planes now have an AED on board. The guidelines recommend that this equipment should be available on all planes.
- It is very important for the crew to request help as soon as possible by an onboard announcement after identification of a patient with cardiac arrest. The announcement should state there has been a suspected cardiac <u>arrest</u> and also the location of the <u>emergency</u> equipment.



- Two-person CPR is considered optimum (as it is better quality) and should be performed if possible; the crew should be trained regularly in basic life support, ideally with a focus on CPR in aircraft.
- The plane should be diverted immediately if the patient has a return of spontaneous circulation.

Dr Hinkelbein says: "This is the first guideline providing specific treatment recommendations for in-flight medical emergencies during commercial air travel. This is of major importance to recommend proper actions and procedures since the airplane environment as well as equipment will be significantly different to what can be provided for medical emergencies on the ground."

He and his team will now begin raising awareness of the guideline by directly contacting individual airlines and asking them to incorporate it into their emergency procedures.

## Provided by European Society of Anaesthesiology

Citation: Cardiac arrest on plane journeys: New guidelines proposed (2017, June 5) retrieved 11 May 2024 from <u>https://medicalxpress.com/news/2017-06-cardiac-plane-journeys-guidelines.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.