

Sleep duration impacts treatment response for depressed patients with insomnia

June 5 2017



Credit: CC0 Public Domain

Preliminary results from a new study show that depressed patients with insomnia who sleep seven or more hours per night are more likely to benefit from cognitive behavioral therapy for insomnia (CBTI) and

achieve depression remission.

Results show that when [insomnia](#) and [depression](#) co-occur, longer pre-treatment objective sleep duration is predictive of remission of both disorders when patients are given a combination of CBTI for insomnia and antidepressant medication for depression.

"A seven-hour, objective sleep duration of patients prior to entering treatment increased their chances of achieving both depression and insomnia remission by their treatment endpoints," said lead author and co-principal investigator Jack D. Edinger, PhD, professor in the Section of Sleep Medicine at National Jewish Health in Denver, Colorado.

The study involved 104 adults, including 75 women, who enrolled in the Treatment of Insomnia and Depression Study and completed one baseline night of polysomnography. Participants received 16 weeks of anti-depressant medication and were randomly assigned either to CBTI or sham insomnia therapy. The Hamilton Rating Scale for Depression (HAM-D-17) and Insomnia Severity Index were administered at baseline and then bi-weekly during treatment to determine depression and insomnia remission.

The study was part of a larger research project for which Edinger was co-investigator along with co-investigators Daniel Buysse, MD, from the University of Pittsburgh; Andy Krystal, MD, from Duke University and the University of California, San Francisco; and lead principal investigator Rachel Manber, PhD, professor of psychiatry and behavioral sciences at the Stanford University Medical Center.

"Our findings highlight the importance of adequate objective sleep in the recovery from depression and insomnia," said Manber. "The data suggest that short sleep duration may be a risk for refractory depression."

The research abstract was published recently in an [online supplement of the journal *Sleep*](#) and will be presented Monday, June 5, in Boston at SLEEP 2017, the 31st Annual Meeting of the Associated Professional Sleep Societies LLC (APSS), which is a joint venture of the American Academy of Sleep Medicine and the Sleep Research Society.

More information: Abstract Title: Does Objective Sleep Duration Moderate Treatment Response in Patient with Comorbid Depression and Insomnia? A report from the TRIAD study

Abstract ID: 0335

Presentation Date: Monday, June 5

Oral Presentation: 2:30 p.m. to 2:45 p.m., ballroom C

Presenter: Jack Edinger, PhD

Provided by American Academy of Sleep Medicine

Citation: Sleep duration impacts treatment response for depressed patients with insomnia (2017, June 5) retrieved 6 May 2024 from <https://medicalxpress.com/news/2017-06-duration-impacts-treatment-response-depressed.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--