

Fatality rates are high within two years of drug-induced liver injury

June 26 2017

Most patients suffering liver toxicity due to medications or herbal or dietary supplements recover from the acute liver injury without long-term problems, but some do not survive the injury or they require liver transplantation. In a recent study, nearly 10% of patients with definite, highly likely, or probable drug-induced liver injury had a fatal outcome within 2 years of onset.

In the 107 deaths among 1089 patients in the study, drug-induced liver injury was the primary cause of death in 68 patients and was a contributing cause in another 15 (at total of 7.6%). The remaining 22 patients had fatal outcomes unrelated to drug-induced liver injury.

The findings are published in *Hepatology*.

More information: Paul H. Hayashi et al, Death and Liver Transplantation within Two Years of Onset of Drug-Induced Liver Injury, *Hepatology* (2017). [DOI: 10.1002/hep.29283](https://doi.org/10.1002/hep.29283)

Provided by Wiley

Citation: Fatality rates are high within two years of drug-induced liver injury (2017, June 26) retrieved 10 April 2024 from <https://medicalxpress.com/news/2017-06-fatality-high-years-drug-induced-liver.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.