


HIV testing inadequate in young men who have sex with men

June 26 2017



(HealthDay)—Few young men who have sex with men (MSM) receive

HIV testing, although they account for most new diagnoses, according to research published in the June 23 issue of the U.S. Centers for Disease Control and Prevention's *Morbidity and Mortality Weekly Report*.

Noting that the CDC recommends screening all persons aged 13 to 64 years for HIV infection, Renee Stein, Ph.D., from the National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention at the CDC in Atlanta, and colleagues analyzed data for [young people](#) submitted by 61 health departments in 2015.

The researchers found that young MSM received 28 percent of HIV tests, although they accounted for 83 percent of new diagnoses. Sixty-six percent of young people who received positive test results for HIV infection were linked to care within 90 days of diagnosis in this analysis.

"Increasing the number of youths at risk for HIV infection who are tested for HIV on a regular basis and ensuring that youths who receive positive test results for HIV are rapidly linked to and retained in appropriate medical care, including early initiation of antiretroviral therapy, are essential steps for reducing HIV infection in this vulnerable population," the authors write.

More information: [Abstract/Full Text](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: HIV testing inadequate in young men who have sex with men (2017, June 26) retrieved 23 April 2024 from <https://medicalxpress.com/news/2017-06-hiv-inadequate-young-men-sex.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.