

Human reproduction likely to be more efficient than previously thought

June 13 2017



Dos rayitas. Credit: Esparta Palma

How difficult is it to conceive? According to a widely-held view, fewer than one in three embryos make it to term, but a new study from a researcher at the University of Cambridge suggests that human embryos are not as susceptible to dying in the first weeks after fertilisation as often claimed.

Dr Gavin Jarvis from Cambridge's Department of Physiology,

Development and Neuroscience re-examined data going back to the 1940's and concluded that previous claims about natural embryo mortality are too often exaggerated. His report is published in *F1000Research*.

"Trying to determine whether a human embryo survives during the first days after fertilisation is almost impossible," says Dr Jarvis. "A woman can only suspect that she is pregnant, at the earliest, two weeks after fertilisation, when she misses a period. Using sensitive laboratory tests, embryos can be detected as they implant into the womb about one week after fertilisation. What happens before then under natural circumstances is anyone's guess."

In 1938, two doctors in Boston, Dr Arthur Hertig and Dr John Rock, became the first people to see a human embryo when they examined wombs removed from women during surgery. They estimated that a half of [human embryos](#) die in the first two weeks after fertilisation. However, Dr Jarvis's re-analysis of this data shows that this figure is so imprecise as to be of little value.

"I think it is fair to say that their data show that embryos can and do fail at these early stages, and also that many do just fine, but we could say that even without the data," he adds. "Hertig's samples, whilst descriptively informative, are quantitatively unhelpful. It doesn't take us much further than where we would be without the data."

Pregnancies are also lost after the first two weeks and currently published estimates of total embryo loss from fertilisation through to birth range from less than 50% to 90%. Embryo mortality of 90% implies that only 10% of embryos survive to birth, implying that human reproduction is highly inefficient.

Since 1988, several studies on women trying to get pregnant have

provided a more consistent picture. The earliest point at which pregnancy can be detected is one week after fertilisation when the embryo starts to implant into the womb of the mother. At this point the hormone hCG, which is used in regular pregnancy tests, becomes detectable. Among implanting embryos, about one in five fail very soon and the woman will have a period at about the expected time, never suspecting that she conceived. Once a period is missed and pregnancy confirmed, about 10-15% will be lost before live birth, mostly within the first few months. In total, once implantation starts, about two thirds of embryos survive to birth. The number of embryos that survive and die before implantation remains unknown.

Modern reproductive technologies have enabled [fertilisation](#) to be observed directly in the laboratory. Poor survival of in vitro embryos may have contributed to the pessimistic view about natural human embryo survival, says Dr Jarvis.

"Fertilising human eggs and culturing human embryos in the laboratory is not easy. A large proportion of eggs fertilised in vitro do not develop properly even for a week. Of those that do and are transferred into women undergoing IVF treatment, most do not become a new-born baby."

This failure of in vitro embryos may reflect the natural situation. Alternatively, the artificial environment of reproductive treatments may contribute to the high failure rate of IVF embryos. Dr Jarvis's re-analysis of the data suggests that the latter is the case.

"It's impossible to give a precise figure for how many [embryos](#) survive in the first week but in normal healthy women, it probably lies somewhere between 60-90%. This wide range reflects the lack of relevant data. Although we can't be precise, we can avoid exaggeration, and from reviewing the studies that do exist, it is clear that many more survive

than is often claimed," concludes Dr Jarvis.

More information: Gavin E. Jarvis. Early embryo mortality in natural human reproduction: What the data say, *F1000Research* (2017). [DOI: 10.12688/f1000research.8937.2](https://doi.org/10.12688/f1000research.8937.2)

Provided by University of Cambridge

Citation: Human reproduction likely to be more efficient than previously thought (2017, June 13) retrieved 20 April 2024 from <https://medicalxpress.com/news/2017-06-human-reproduction-efficient-previously-thought.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.