


## Keep well hydrated to help keep kidney stones away

June 13 2017, by Kristi Lopez



Credit: CC0 Public Domain

Each year, more than a million people in the U.S. will seek treatment for mild to severe pain caused by a kidney stone. Overall, one in 11 individuals in the U.S. will be affected by kidney stones at some time in their life. But one of the most common times to form kidney stones is during the summer and autumn months.



Kidney stones are hard, pebble-like pieces of material that form in one or both kidneys when your <u>urine</u> contains more crystal-forming substances—such as calcium, oxalate and uric acid—than the fluid in your urine can dilute. In addition, your urine may lack substances that prevent particles from sticking together, creating an ideal environment for kidney stones to form.

Kidney stones can vary in size from tiny crystals that can only be seen with a microscope to stones more than an inch wide. Tiny stones may pass without your even noticing but can be extremely painful. Stones of any size can become stuck in the <u>urinary tract</u> causing swelling of the kidney and intractable flank pain.

The most effective way to prevent kidney stones from forming is to drink plenty of water. Dehydration is one of the most common reasons why people form stones. When we are dehydrated, urine is too concentrated and minerals can build up and form stones.

In fact, during the warmer or hotter times of the year, you're at greatest risk of becoming dehydrated and even more important to drink more than you usually drink even during the cooler days or months.

Kidney stones can form at any age, but they usually appear in those between 40 and 60 years old. Of those who develop one stone, half will develop at least one more in the future.

Symptoms may not appear until the stone moves around within the kidney or passes into the ureter – the tube connecting the kidney and bladder. The most common signs and symptoms are:

- Severe pain in the side and back, below the ribs
- Pain that radiates to the lower abdomen and groin
- Pain that comes in waves and fluctuates in intensity



- Pain on urination
- Pink, red or brown urine
- Cloudy or foul-smelling urine
- Nausea and vomiting
- Persistent need to urinate
- Urinating more often than usual
- Fever and chills if an infection is present
- Urinating small amounts

If you have these signs or symptoms, seek medical attention. Treatment depends on the stone's size and location and a CT scan or X-ray can help pinpoint the location and estimate the size of a kidney stone. Depending on what your doctor finds, you may be prescribed medicine and advised to drink a lot of fluids. Or, you might need a procedure to break up or remove the kidney stone.

Once you've had a <u>kidney stone</u>, you have an increased chance for having another. So, as you head outdoors this summer, don't forget to hydrate.

Provided by University of Kentucky

Citation: Keep well hydrated to help keep kidney stones away (2017, June 13) retrieved 28 April 2024 from <u>https://medicalxpress.com/news/2017-06-hydrated-kidney-stones.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.