

New insights into exercise right ventricular pressure may help define a new 'normal'

June 21 2017

In individuals with structurally normal hearts, systolic pressure is assumed to be equal in the lung arteries and the part of the heart pumping to them. A difference in pressure between the right ventricle (RV) and pulmonary artery (PA) suggests an obstruction to blood flow (specifically in the right ventricular outflow tract, or RVOT), which is most often seen in patients who have congenital heart defects or sometimes after cardiac surgery or lung transplantation. An RVOT pressure gradient is usually seen as a sign for concern: if the obstruction to the blood flow is severe, it can lead to right heart failure.

However, investigators at Brigham and Women's Hospital (BWH) recently reported an unexpected observation among patients experiencing unexplained shortness of breath. Their findings, reported in *PLOS One*, suggest that the RV-PA pressure difference may not always be a reflection of disease, but rather, may be a normal physiological response to <u>exercise</u>.

The research team observed that when patients underwent cardiopulmonary exercise testing, a subset of them (who had normal resting RV-PA pressure) developed the pressure gradient during high levels of upright exercise when the <u>heart</u> was squeezing. In a retrospective study, the investigators reviewed data from patients referred to the BWH Dyspnea Center who underwent an invasive, comprehensive exercise test with a catheter in place to measure RV and PA pressures. Their body's response was measured through heart catheterization when the patients were resting supine (lying down), but



also when they were in an upright position, exercising on a stationary bike until their ability to keep exercising was limited by symptoms. Usually, patients are only tested in the supine position at rest; testing upright patients provided new information about when and in which patients an RVOT pressure gradient may occur.

Surprisingly, those in the study who developed the RVOT pressure gradient during high levels of upright exercise were not more limited. Rather, they tended to be the younger <u>patients</u> with better exercise capacity.

"Our work suggests that, surprisingly, this pressure gradient during exercise may represent a normal, physiologic state in fit, young people," said corresponding author Alexander Opotowsky, MD, MPH of the Pulmonary Vascular Disease Team and the Dyspnea Center at BWH and of Boston Adult Congenital Heart Service, a joint program of Boston Children's Hospital and BWH.

Due to the scope of this study, additional testing among asymptomatic, young, aerobically fit individuals will need to be conducted in order to confirm the suggestive findings and better understand the RVOT gradient. The reasoning behind why an RVOT gradient would form during upright exercise currently remains unclear and further studies to illuminate the mechanism responsible for the observed RVOT gradient are needed. But importantly, the results of this study may have implications for the standard practice of exercise echocardiography.

"Currently, some of our testing with echocardiography depends on the fact that the RV and PA have the same pressure," said Opotowsky. "Our findings raise important questions about this assumption and may have implications for how echocardiography is used to estimate systolic <u>pulmonary artery pressure</u> during exercise."



More information: van Riel AC et al. "Hemodynamic and metabolic characteristics associated with development of a right ventricular outflow tract pressure gradient during upright exercise" *PLOS One* (2017). DOI: 10.1371/journal.pone.0179053

Provided by Brigham and Women's Hospital

Citation: New insights into exercise right ventricular pressure may help define a new 'normal' (2017, June 21) retrieved 3 May 2024 from <u>https://medicalxpress.com/news/2017-06-insights-ventricular-pressure.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.