

Review links conjugated linoleic acid supplementation to CRP

June 24 2017



(HealthDay)—Conjugated linoleic acid (CLA) supplementation is



associated with an increase in serum C-reactive protein (CRP) concentration, according to a review and meta-analysis published online May 29 in *Cardiovascular Therapeutics*.

Mohsen Mazidi, Ph.D., from the Chinese Academy of Sciences in Beijing, and colleagues conducted a systematic review and meta-analysis to examine the impact of CLA supplementation on serum CRP. Data were included from 14 studies.

The researchers found that following supplementation with CLAs there was a significant increase in serum CRP concentrations (weighted mean difference, 0.63 mg/dL; 95 percent confidence interval, 0.13 to 1.13; heterogeneity P = 0.026); these findings were robust in sensitivity analyses. The changes in serum CRP levels were found to be independent of CLA supplementation dosage (slope, -0.02; 95 percent confidence interval, -0.10 to 0.12; P = 0.889) or follow-up duration (slope, 0.271; 95 percent confidence interval, -0.05 to 0.59; P = 0.098).

"This meta-analysis suggests that CLA supplementation is associated with an increase in plasma CRP concentrations and a reduction in <u>serum</u> adiponectin concentrations, which indicates that CLA <u>supplements</u> have a pro-inflammatory effect," the authors write.

More information: Abstract

Full Text (subscription or payment may be required)

Copyright © 2017 HealthDay. All rights reserved.

Citation: Review links conjugated linoleic acid supplementation to CRP (2017, June 24) retrieved 9 April 2024 from

https://medicalxpress.com/news/2017-06-links-conjugated-linoleic-acid-supplementation.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.