


# Longer survival for midostaurin and chemotherapy in AML with FLT3

June 26 2017

---



(HealthDay)—For patients with acute myeloid leukemia (AML) and a

*FLT3* mutation, midostaurin plus chemotherapy is associated with prolonged overall and event-free survival, according to a study published online June 23 in the *New England Journal of Medicine*.

Richard M. Stone, M.D., from the Dana-Farber Cancer Institute in Boston, and colleagues screened 3,277 [patients](#) aged 18 to 59 years with newly diagnosed AML for *FLT3* mutations. Patients were randomized to receive either standard chemotherapy plus midostaurin or placebo; those who were in remission after consolidation therapy entered a maintenance phase with midostaurin or placebo. Randomization was stratified according to *FLT3* mutation subtype.

Three hundred sixty patients were randomized to the midostaurin group and 357 to the [placebo group](#). The researchers found that overall survival was significantly longer in the midostaurin versus the [placebo](#) group (hazard ratio for death, 0.78), as was event-free survival (hazard ratio for event or death, 0.78). The benefit of midostaurin was consistent across all *FLT3* subgroups in the primary analysis and in an analysis in which data for patients who underwent transplantation were censored.

"The addition of the multitargeted kinase inhibitor midostaurin to standard chemotherapy significantly prolonged overall and event-free survival among patients with AML and a *FLT3* mutation," the authors write.

The study was partially funded by Novartis, the manufacturer of midostaurin.

**More information:** [Abstract](#)  
[Full Text](#)

Copyright © 2017 [HealthDay](#). All rights reserved.

Citation: Longer survival for midostaurin and chemotherapy in AML with FLT3 (2017, June 26)  
retrieved 25 April 2024 from

<https://medicalxpress.com/news/2017-06-longer-survival-midostaurin-chemotherapy-aml.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.